

## Three-Dimensional Orthogonal Graph Drawing with Optimal Volume<sup>1</sup>

Therese Biedl,<sup>2</sup> Torsten Thiele,<sup>3</sup> and David R. Wood<sup>4</sup>

**Abstract.** An orthogonal drawing of a graph is an embedding of the graph in the rectangular grid, with vertices represented by axis-aligned boxes, and edges represented by paths in the grid that only possibly intersect at common endpoints. In this paper we study three-dimensional orthogonal drawings and provide lower bounds for three scenarios: (1) drawings where the vertices have a bounded aspect ratio, (2) drawings where the surfaces of vertices are proportional to their degrees, and (3) drawings without any such restrictions. Then we show that these lower bounds are asymptotically optimal, by providing constructions that in all scenarios match the lower bounds within a constant factor.

**Key Words.** Graph drawing, Orthogonal, Three-dimensional, Lower bounds.

**1. Introduction.** Graph drawing is a field with a wide range of applications, for example in network visualisation, data base design, and telecommunications. See the recent book [11] for an overview of techniques in graph drawing. Orthogonal graph drawing, where edges are routed along a rectangular grid, is a popular drawing style with applications in data flow diagrams, entity relationship diagrams, and VLSI design.

In this paper we study three-dimensional orthogonal graph drawings, and in particular, focus on minimising the volume of such drawings. We improve on previous results by generalising the existing lower bounds on the volume, and giving new constructions with smaller volume. In fact, for three different drawing scenarios, our upper and lower bounds are matching up to a constant factor, and are hence asymptotically optimal. To state our results precisely, we first give formal definitions and notation.

The (*three-dimensional*) *rectangular grid* is the cubic lattice, consisting of *grid points* with integer coordinates, together with the axis-parallel *grid lines* determined by these points. We use the word *box* to mean a three-dimensional axis-parallel box whose corners have integer coordinates. That is, a box  $B$  is a set of points

$$\{(p_x, p_y, p_z) : l_I \leq p_I \leq r_I, I \in \{X, Y, Z\}\}$$

<sup>1</sup> A preliminary version of this paper was presented at the 8th International Symposium on Graph Drawing (GD 2000), Colonial Williamsburg, Virginia, USA, September 20–23, 2000.

<sup>2</sup> School of Computer Science, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1. [biedl@uwaterloo.ca](mailto:biedl@uwaterloo.ca). Research supported by NSERC.

<sup>3</sup> Institut für Informatik, Freie Universität Berlin, Takustraße 19, 14195 Berlin, Germany. [thiele@inf.fu-berlin.de](mailto:thiele@inf.fu-berlin.de).

<sup>4</sup> Department of Mathematics and Statistics, and School of Computer Science, McGill University, Montréal, Quebec, Canada H3A 2A7. [wood@cs.mcgill.ca](mailto:wood@cs.mcgill.ca). Research completed at Monash University (Melbourne, Australia), The University of Sydney (Sydney, Australia), and Carleton University (Ottawa, Canada).

Received January 23, 2003; revised March 23, 2004, and December 6, 2004. Communicated by T. Nishizeki. Online publication May 2, 2005.

for some integers  $l_I, r_I, I \in \{X, Y, Z\}$ . At each grid point in a box  $B$  that is extremal in some direction  $d \in \{\pm X, \pm Y \pm Z\}$ , there is *port* on  $B$  in direction  $d$ . One grid point can thus define up to six incident ports. For each dimension  $I \in \{X, Y, Z\}$ , an  $I$ -line is a line parallel to the  $I$ -axis, an  $I$ -segment is a line-segment within an  $I$ -line, and an  $I$ -plane is a plane perpendicular to the  $I$ -axis.

Let  $G = (V, E)$  be a graph, which is allowed to have parallel edges but no self-loops. We denote the number of vertices of  $G$  by  $n = |V|$ , the number of edges of  $G$  by  $m = |E|$ , and the maximum degree of  $G$  by  $\Delta(G)$ , or  $\Delta$  if the graph in question is clear.

An *orthogonal (box-)drawing* of  $G$  represents vertices by pairwise non-intersecting boxes. Hence vertices are possibly degenerate, in the sense that they may be represented by a rectangle or even a line-segment or a point. This is the approach taken in [5], [8], [25], and [26], but not in [21]. (Enlarging vertices to remove this degeneracy increases the volume by a multiplicative constant.) An edge  $vw \in E$  is represented by a sequence of contiguous segments of grid lines possibly bent at grid points, between ports on the boxes of  $v$  and  $w$ . The intermediate grid points along the path representing an edge do not intersect the box of any vertex or any other edge route.

An orthogonal drawing with a particular shape of box representing every vertex, e.g. point, line-segment, or cube, is called an orthogonal *shape-drawing* for each particular shape. Initial research in orthogonal drawing was mostly concerned with point-drawings; see for example [9], [10], [12], [13], [21], [28], and [29]. However, three-dimensional orthogonal point-drawings can only exist for graphs with maximum degree at most six. Overcoming this restriction has motivated recent interest in orthogonal box-drawings [5], [8], [21], [25], [26]. Related research in three-dimensional VLSI includes [1], [2], [18], [22], and [23].

From now on, we use the term *drawing* to mean a three-dimensional orthogonal box-drawing. Furthermore, the graph-theoretic terms “vertex” and “edge” also refer to their representation in a drawing. The *size* of a vertex  $v$  in a drawing is denoted by  $X(v) \times Y(v) \times Z(v)$ , where for each  $I \in \{X, Y, Z\}$ ,  $I(v)$  is one more than the length of the side of (the box of)  $v$  parallel to the  $I$ -axis. Thus, if the boundaries of  $v$  are  $r_I(v)$  and  $l_I(v)$  then  $I(v) = r_I(v) - l_I(v) + 1$ . The number of ports of  $v$  is called its *surface*, denoted by  $\text{surface}(v)$ . The number of grid points in a box is called its *volume*.

The smallest box enclosing a drawing is called the *bounding box* of the drawing. The *volume* of a drawing is the volume of its bounding box. The volume and the maximum number of bends per edge are the most commonly proposed measures for determining the quality of a drawing. For box-drawings, the size and shape of a vertex with respect to its degree are also considered an important measure of quality. We use the following two measures for the shape of a vertex:

**DEGREE-RESTRICTED DRAWINGS.** A vertex  $v$  in a drawing is *strictly  $\alpha$ -degree-restricted* if

$$\text{surface}(v) \leq \alpha \cdot \text{deg}(v)$$

for some constant  $\alpha \geq 1$ . If there exists a constant  $\alpha$  such that every vertex in a drawing is strictly  $\alpha$ -degree-restricted then the drawing is *strictly degree-restricted*, or *strictly  $\alpha$ -degree-restricted* if we wish to emphasise the value of the constant.

For some drawing algorithms, the minimum  $\alpha$  such that the drawings produced by the algorithm are strictly  $\alpha$ -degree-restricted does not necessarily reflect the asymptotic relationship between the surface and the degree of the vertices. We therefore say that in a drawing, a vertex  $v$  is  $\alpha$ -degree-restricted if

$$\text{surface}(v) \leq \alpha \cdot \deg(v) + o(\deg(v)).$$

If for some constant  $\alpha$ , every vertex  $v$  is  $\alpha$ -degree-restricted then the drawing is *degree-restricted* or  $\alpha$ -*degree-restricted*. This definition enables us to compare the asymptotic behaviour of  $\alpha$  for various algorithms.

Clearly, if a drawing is strictly degree-restricted then it is also degree-restricted. Conversely, it is easily seen that the degree-restricted drawings produced by the algorithm presented in this paper (and all known algorithms) are also strictly degree-restricted, thus for our purposes the two notions coincide. It is necessary to distinguish the two terms as the lower bound in Theorem 2 is for strictly degree-restricted drawings.

**ASPECT RATIO.** The *aspect ratio* of a vertex  $v$  is the ratio between its largest and smallest side, i.e.,

$$\max\{X(v), Y(v), Z(v)\} / \min\{X(v), Y(v), Z(v)\}.$$

In particular, a vertex with aspect ratio one is a cube. A drawing *has bounded aspect ratios* if there exists a constant  $r$  such that all vertices have aspect ratios at most  $r$ . Note that necessarily  $r \geq 1$ .

There is no inherent relationship between whether a drawing is degree-restricted or has bounded aspect ratios. Previously, algorithms have been presented that give drawings that are degree-restricted, but do not have bounded aspect ratios [5], [25], [26]. It is conceivable that a drawing could have bounded aspect ratios, but not be degree-restricted (for example, by representing each vertex  $v$  with a  $\deg(v) \times \deg(v) \times \deg(v)$  box), though no algorithms to create such drawings have been presented, and as our lower bound results show, no improvement in volume is possible by doing so.

**1.1. Lower Bounds.** For a graph  $G$ , denote by  $\text{vol}(G, r, \alpha)$  the minimum volume, taken over all (orthogonal) drawings of  $G$  that have aspect ratios at most  $r$  and are strictly  $\alpha$ -degree-restricted. Let  $\text{vol}(n, m, r, \alpha)$  be the maximum, taken over all graphs  $G$  with  $n$  vertices and  $m$  edges, of  $\text{vol}(G, r, \alpha)$ . Thus,  $\text{vol}(n, m, r, \alpha)$  describes a volume bound within which we can draw all graphs with  $n$  vertices and  $m$  edges such that each vertex  $v$  has aspect ratio at most  $r$  and surface at most  $\alpha \cdot \deg(v)$ .

The first lower bounds on the volume were due to Hagihara et al. [16].<sup>5</sup> They show that, in the above notation,  $\text{vol}(n, m, 1, 1) = \Omega(\max\{n\Delta^2, (n\Delta/\log n)^{3/2}\})$ . In fact, in their construction the graphs are  $\Delta$ -regular. Hence  $m = \frac{1}{2}n\Delta$ , which allows us to restate their result as

$$\text{vol}(n, m, 1, 1) = \Omega(\max\{m\Delta, (m/\log n)^{3/2}\}).$$

<sup>5</sup> This paper seems widely unknown in the graph drawing community.

In this paper, we show that:

- $vol(n, m, \infty, \infty) = \Omega(m\sqrt{n})$ .
- $vol(n, m, r, \infty) = \Omega(m^{3/2}/\sqrt{r})$ .
- $vol(n, m, \infty, \alpha) = \Omega(m^{3/2}/\alpha)$ .

We thus improve the results of [16] in three ways: Firstly, we remove the log-factor to establish  $vol(n, m, 1, 1) = \Omega(m^{3/2})$  as the lower bound. Secondly, we show that the result holds even if only one of the conditions of having bounded aspect ratios and being strictly degree-restricted holds. Finally, we also study the case when neither of these two conditions hold. The lower bound here appears weaker, but as we prove by giving a construction later, it is asymptotically optimal.

Our first result includes the lower bound of  $\Omega(n^{5/2})$  for drawings of  $K_n$  established by Biedl et al. [8]. In fact, the proofs of our lower bounds are based on techniques developed in that paper, generalised to graphs with fewer edges.

*1.2. Algorithms.* A trade-off between the maximum number of bends per edge route and the bounding box volume is apparent in algorithms for orthogonal graph drawing. (Eades et al. [13] discuss this phenomena in the case of orthogonal point-drawings.) Biedl et al. [8] construct drawings of  $K_n$  with  $\mathcal{O}(n^{5/2})$  volume and three bends per edge, but these drawings are not degree-restricted. They also construct drawings of  $K_n$  with  $\mathcal{O}(n^3)$  volume and one bend per edge, but these drawings are degree-restricted only for graphs where all vertices have degree  $\theta(n)$ . Wood [27] obtained drawings with one bend per edge and  $\mathcal{O}(n^{3/2}m)$  volume. Biedl [4] showed that  $\Omega(n^3)$  volume is required for  $K_n$  if only one bend per edge is allowed.

A drawing is in *general position* if no two vertices are in a common grid plane. The algorithm of Papakostas and Tollis [21] produces general position drawings with  $\mathcal{O}(m^3)$  volume. This bound has been improved to  $\mathcal{O}((nm)^{3/2})$  for cube-drawings and  $\mathcal{O}(n^2m)$  for line-segment-drawings in general position by Biedl [5] and Wood [25], respectively.

The *lifting half-edges* technique developed by Biedl [5] generates drawings of simple graphs starting with a two-dimensional general position drawing [7] (possibly with overlapping edges). The edge routes are partitioned into sub-drawings each consisting of  $X$ -segments or  $Y$ -segments. Each sub-drawing is then assigned its own  $Z$ -plane, vertices are extended to form lines passing through each layer, and vertical segments are added to the edges in such a way as to avoid crossings. Using this technique an improved volume bound of  $\mathcal{O}(n^2\Delta)$  is attained. This bound has been improved to  $\mathcal{O}(nm)$ , although the drawings are no longer degree-restricted [27]. At the cost of an increase in volume, cube-drawings can also be produced in the lifting half-edges model.

The *plane-drawing* technique consists of positioning vertices in the  $(Z = 0)$ -plane, and then routing edges above and possibly also below the  $(Z = 0)$ -plane. The two algorithms presented in this paper use the plane-drawing technique. The first algorithm produces degree-restricted cube-drawings with  $\mathcal{O}(m^{3/2})$  volume and at most six bends per edge. The technique used is a generalisation of the COMPACT algorithm of Eades et al. [13] for point-drawings, and is an improvement on the algorithms of Hagihara et al. [16] and Wood [26], who obtained upper bounds of  $\mathcal{O}((n\Delta)^{3/2})$  and  $\mathcal{O}(m^2/\sqrt{n})$ , respectively. Our second algorithm produces box-drawings with  $\mathcal{O}(m\sqrt{n})$  volume and at most four

**Table 1.** The trade-off between volume and the maximum number of bends in orthogonal graph drawings for various quality measures. All lower bounds are proved in Theorem 2.

Lower Bound	Volume	Bends	Model	Graphs	Reference
Bounded aspect ratio / degree-restricted					
$\Omega(m^{3/2})$	$\mathcal{O}((nm)^{3/2})$	2	General position	Simple	[5], [25]
$\Omega(m^{3/2})$	$\mathcal{O}(nm\sqrt{\Delta})$	2	Lifting half-edges	Simple	[5]
$\Omega(m^{3/2})$	$\mathcal{O}(m^2)$	5	Plane layout	Multigraphs	Theorem 4
$\Omega(m^{3/2})$	$\mathcal{O}((n\Delta)^{3/2})$	10	Plane layout	Simple	[16]*
$\Omega(m^{3/2})$	$\mathcal{O}(m^{3/2})$	6	Plane layout	Multigraphs	Theorem 3
No bounds on aspect ratio / degree-restricted					
$\Omega(m^{3/2})$	$\mathcal{O}(n^2m)$	2	General position	Simple	[5], [25]
$\Omega(m^{3/2})$	$\mathcal{O}(n^2\Delta)$	2	Lifting half-edges	Simple	[5]
$\Omega(m^{3/2})$	$\mathcal{O}(m^2)$	5	Plane layout	Multigraphs	Theorem 4
$\Omega(m^{3/2})$	$\mathcal{O}(m^{3/2})$	6	Plane layout	Multigraphs	Theorem 3
No bounds on aspect ratio / not necessarily degree-restricted					
$\Omega(m\sqrt{n})$	$\mathcal{O}(n^3)$	1	Lifting edges	Simple	[8]
$\Omega(m\sqrt{n})$	$\mathcal{O}(n^{3/2}m)$	1	Lifting edges	Simple	[27]
$\Omega(m\sqrt{n})$	$\mathcal{O}(nm)$	2	Lifting half-edges	Simple	[27]
$\Omega(m\sqrt{n})$	$\mathcal{O}(n^{5/2})$	3	Lifting edges	Simple	[8]
$\Omega(m\sqrt{n})$	$\mathcal{O}(nm)$	3	Plane layout	Multigraphs	Theorem 6
$\Omega(m\sqrt{n})$	$\mathcal{O}(m\sqrt{n})$	4	Plane layout	Simple	Theorem 5

\*Hagihara et al. [16] did not consider the number of bends per edge; we deduce the bound of 10 from their construction.

bends per edge; these drawings are not degree-restricted nor do the vertices have bounded aspect ratios.<sup>6</sup>

Both upper bounds are therefore within a constant factor of the lower bound. We also present refinements of both our algorithms with one less bend per edge, at the cost of an increase in the volume. Table 1 summarises the known bounds for orthogonal graph drawing.

**2. Lower Bounds.** In this section we prove lower bounds on the volume of orthogonal graph drawings. Such lower bounds were previously only known for drawings of the complete graph  $K_n$  [8]. The crucial argument for  $K_n$  is that between any two disjoint vertex sets of size  $\theta(n)$  in  $K_n$ , there are  $\theta(n)^2$  edges. To generalise this to arbitrary graphs, we first exhibit graphs such that between any two disjoint vertex sets of size  $\theta(n)$  there are  $\theta(m)$  edges. Then we use an argument similar to that in [8] to obtain lower bounds on the volume.

**2.1. Graphs with Large Cuts.** Suppose  $G = (V, E)$  is a graph and  $S, T \subset V$  are disjoint sets of vertices. Let  $e(S, T)$  denote the number of edges between  $S$  and  $T$ . For our lower bound proofs, we need graphs for which  $e(S, T)$  is large under some conditions on  $S$  and  $T$ , and we prove the existence of such graphs in the following lemma.

<sup>6</sup> These results assume that  $m \geq n$ , which one would expect in most applications. If  $m < n$  then the volume of the drawings produced is  $\mathcal{O}(n^{3/2})$  in both constructions.

LEMMA 1. *If  $p \neq q$  are primes,  $p \equiv 1 \pmod{4}$ ,  $q \equiv 1 \pmod{4}$ ,  $144 \leq p < q(q-1)/2$ , then there exists a simple  $n$ -vertex graph  $G_{p,q} = (V, E)$  with the following properties:*

- $G_{p,q}$  is  $d$ -regular with  $d = p + 1$ .
- $q(q-1)/2 \leq n \leq q(q-1)$ .
- For any disjoint sets  $S, T \subset V$  of vertices of  $G_{p,q}$  with  $|S||T| \geq n^2/36$  we have

$$e(S, T) \geq C \cdot dn,$$

where  $C \approx 0.00009$  is a constant.

Before proving this lemma, we need some background. For a graph  $G$ , denote by  $\lambda(G)$  the second largest eigenvalue of its adjacency matrix. The following well-known inequality (see, for example, [3, pp. 119–125] and [24]) relates  $\lambda(G)$  to the cut property we are interested in.

LEMMA 2. *Let  $G = (V, E)$  be a  $d$ -regular  $n$ -vertex graph with second largest eigenvalue  $\lambda(G)$ . Then, for all disjoint sets  $S, T \subset V$ , we have*

$$e(S, T) \geq \frac{d|S||T|}{n} - \lambda(G)\sqrt{|S||T|}.$$

This lemma suggest looking for graphs  $G$  with small  $\lambda(G)$ . Fortunately, such graphs (called *Ramanujan graphs*) have already been constructed.

LEMMA 3 [19], [20]. *If  $p \neq q$  are primes,  $p \equiv 1 \pmod{4}$ ,  $q \equiv 1 \pmod{4}$ , then there exists a simple  $n$ -vertex graph  $G_{p,q} = (V, E)$  with the following properties:*

- $G_{p,q}$  is  $d$ -regular with  $d = p + 1$ .
- $q(q-1)/2 \leq n \leq q(q-1)$ .
- $\lambda(G_{p,q}) \leq 2\sqrt{d-1}$ .

REMARK. This second-largest eigenvalue is known to be asymptotically optimal, see [19].

Now we are in a position to prove Lemma 1.

PROOF. Given  $p$  and  $q$ , construct the graph  $G_{p,q} = (V, E)$  as in Lemma 3. Let  $S, T \subset V$  be two disjoint sets of vertices of  $G_{p,q}$  with  $|S||T| \geq n^2/36$ . From Lemma 2, we know that

$$e(S, T) \geq \frac{d|S||T|}{n} - \lambda(G_{p,q})\sqrt{|S||T|}.$$

Since  $|S||T| \geq n^2/36$  and  $d = p + 1 \geq 145$ ,

$$\lambda(G_{p,q})\sqrt{|S||T|} \leq 2\sqrt{d-1}\sqrt{|S||T|} \cdot \underbrace{6\sqrt{|S||T|}/n}_{\geq 1} \cdot \underbrace{\sqrt{d}/\sqrt{145}}_{\geq 1} \leq 12/\sqrt{145} \cdot d|S||T|/n.$$

Hence,

$$e(S, T) \geq (1 - 12/\sqrt{145}) \cdot d|S||T|/n \geq \frac{1}{36}(1 - 12/\sqrt{145})dn,$$

which proves the claim for  $C = \frac{1}{36}(1 - 12/\sqrt{145}) \approx 0.000096$ .  $\square$

For future reference, we will call the constant  $C$  the *Ramanujan constant*.

**2.2. Lower Bounds on the Volume of Drawings.** We start by proving the lower bound for those graphs that satisfy the conditions of Lemma 1. The proof is based on the technique developed in [8], which distinguishes three cases: (1) there exists a grid line that intersects many vertices, (2) every grid plane intersects few vertices, or (3) neither of these is the case. Our approach is different in two ways: we use graphs with large cuts, rather than  $K_n$ , and we incorporate considerations of the aspect ratio and degree-restrictions.

**THEOREM 1.** *Let  $G = (V, E)$  be a  $d$ -regular simple graph with  $n \geq 8$  vertices such that for any disjoint sets  $S, T \subset V$  with  $|S||T| \geq n^2/36$  we have  $e(S, T) \geq C \cdot dn$ . Then for all  $\alpha \geq 1$  and for all  $r \geq 1$ ,*

- $\text{vol}(G, \infty, \infty) \geq \frac{1}{3}C^{3/2} \cdot dn^{3/2}$ ,
- $\text{vol}(G, r, \infty) \geq \frac{1}{3}C^{3/2} \cdot (dn)^{3/2}/\sqrt{r}$ ,
- $\text{vol}(G, \infty, \alpha) \geq \frac{1}{3}C^2 \cdot (dn)^{3/2}/\alpha$ .

**PROOF.** Consider an orthogonal drawing of  $G$  in a grid of dimensions  $X \times Y \times Z$ .

*Case 1: A line intersects many vertices.* Assume that there exists a  $Z$ -line that intersects at least  $2\lceil \frac{1}{6}n \rceil$  vertices. Let  $v_1, \dots, v_t$  be the vertices intersected by the  $Z$ -line, listed in order of occurrence along the line. Let  $Z_0$  be a not necessarily integer  $Z$ -coordinate such that the  $(Z = Z_0)$ -plane intersects none of these  $t$  vertices and separates the first  $\lceil \frac{1}{6}n \rceil$  of them from the remaining ones, of which there are at least  $\lceil \frac{1}{6}n \rceil$ .

By assumption at least  $C \cdot dn$  edges connect these two groups. These edges cross the  $(Z = Z_0)$ -plane, which thus must contain at least  $C \cdot dn$  points having integer  $X$ - and  $Y$ -coordinates. Hence  $XY \geq C \cdot dn$ . Since the  $Z$ -line intersects at least  $2\lceil \frac{1}{6}n \rceil \geq \frac{1}{3}n$  vertices, we have  $Z \geq \frac{1}{3}n$ . Thus  $XYZ \geq \frac{1}{3}C \cdot dn^2 \geq \frac{1}{3}C^{3/2} \cdot (dn)^{3/2}$  since  $C \leq 1$  and  $d \leq n$ . This proves all claims.

Similarly, one proves all claims if any  $X$ -line or any  $Y$ -line intersects at least  $2\lceil \frac{1}{6}n \rceil$  vertices.

*Case 2: No plane intersects many vertices.* Assume that any  $X$ -plane,  $Y$ -plane, or  $Z$ -plane intersects at most  $n - 2\lceil \frac{1}{6}n \rceil + 1$  vertices. A vertex is *left* of an  $(X = X_0)$ -plane if all the points in its grid box have  $X$ -coordinates less than  $X_0$ . The notion of *right* of an  $(X = X_0)$ -plane is analogous. As an  $(X = X_0)$ -plane is swept from smaller to larger values of  $X_0$ , it intersects at most  $n - 2\lceil \frac{1}{6}n \rceil + 1$  vertices at any time by assumption.

During the sweep by the  $(X = X_0)$ -plane, an integer  $X'$  is encountered where, for the last time, there are at most  $\lceil \frac{1}{6}n \rceil - 1$  vertices to the left of the  $(X = X')$ -plane. Since the  $(X = X')$ -plane intersects at most  $n - 2\lceil \frac{1}{6}n \rceil + 1$  vertices, there are at least  $\lceil \frac{1}{6}n \rceil$

vertices to the right of the  $(X = X')$ -plane. All these vertices also lie to the right of the  $(X = X' + \frac{1}{2})$ -plane.

By definition of  $X'$ , the number of vertices that lie left of the  $(X = X' + 1)$ -plane is at least  $\lceil \frac{1}{6}n \rceil$ . All these vertices also lie to the left of the  $(X = X' + \frac{1}{2})$ -plane.

By assumption there are at least  $C \cdot dn$  edges between the vertices on the left and the vertices on the right of the  $(X = X' + \frac{1}{2})$ -plane, thus  $YZ \geq C \cdot dn$ . Since the same argument holds for the other two directions,  $XYZ = \sqrt{XY \cdot YZ \cdot XZ} \geq (C \cdot dn)^{3/2}$ , which proves all claims.

*Case 3: A plane intersects many vertices.* Assume now that none of the previous cases hold. Therefore any  $X$ -line,  $Y$ -line, or  $Z$ -line intersects at most  $2\lceil \frac{1}{6}n \rceil - 1$  vertices, but there exists, say, a  $(Z = Z_0)$ -plane that intersects at least  $n - 2\lceil \frac{1}{6}n \rceil + 2 \geq \frac{2}{3}n$  vertices. As an  $(X = X_0)$ -plane is swept from smaller to larger values of  $X_0$ , the  $Y$ -line determined by the intersection of this  $(X = X_0)$ -plane with the  $(Z = Z_0)$ -plane sweeps the  $(Z = Z_0)$ -plane. At any time, this  $Y$ -line intersects at most  $2\lceil \frac{1}{6}n \rceil - 1$  vertices by assumption.

During the sweep by the  $(X = X_0)$ -plane, an integer  $X'$  is encountered where, for the last time, there are at most  $\lceil \frac{1}{6}n \rceil - 2$  vertices left of the  $(X = X')$ -plane and intersecting the  $(Z = Z_0)$ -plane. Since the  $Y$ -line determined by the intersection of the  $(X = X')$ -plane and the  $(Z = Z_0)$ -plane intersects at most  $2\lceil \frac{1}{6}n \rceil - 1$  vertices, and the  $(Z = Z_0)$ -plane intersects at least  $n - 2\lceil \frac{1}{6}n \rceil + 2$  vertices, at least  $n - 5\lceil \frac{1}{6}n \rceil + 5$  vertices intersect the  $(Z = Z_0)$ -plane and lie to the right of the  $(X = X')$ -plane. All these vertices, which we denote by  $S$ , also lie to the right of the  $(X = X' + \frac{1}{2})$ -plane.

By definition of  $X'$ , the number of vertices that intersect the  $(Z = Z_0)$ -plane and that lie left of the  $(X = X' + 1)$ -plane is at least  $\lceil \frac{1}{6}n \rceil - 1$ . All these vertices, which we denote by  $T$ , also lie to the left of the  $(X = X' + \frac{1}{2})$ -plane.

Note that  $|S||T| \geq (n - 5\lceil \frac{1}{6}n \rceil + 5)(\lceil \frac{1}{6}n \rceil - 1)$ , and we claim that this is at least  $\frac{1}{36}n^2$ . If  $n \equiv 0 \pmod{6}$  then  $|S||T| \geq (n - \frac{5}{6}n + 5)(\frac{1}{6}n - 1) = \frac{1}{36}n^2 + \frac{4}{6}n - 5 \geq \frac{1}{36}n^2$  (since  $n \geq 8$ ). If  $n \equiv k \pmod{6}$  for some  $k$ ,  $1 \leq k \leq 5$ , then  $\lceil \frac{1}{6}n \rceil = \frac{1}{6}(n + 6 - k)$  and  $|S||T| \geq (n - \frac{5}{6}(n + 6 - k) + 5)(\frac{1}{6}(n + 6 - k) - 1) = \frac{1}{36}(n^2 + 4kn - 5k^2) \geq \frac{1}{36}n^2$  since  $n \geq 8$  and  $k \leq 5$ .

By assumption there are at least  $C \cdot dn$  edges between  $S$  and  $T$ , thus  $YZ \geq C \cdot dn$ . Apply exactly the same argument in the  $Y$ -direction to obtain  $YZ \geq C \cdot dn$ .

Now we obtain the three lower bounds as follows:

- The  $(Z = Z_0)$ -plane intersects at least  $\frac{2}{3}n$  vertices, thus  $XY \geq \frac{2}{3}n$ . This implies  $XYZ = \sqrt{XY \cdot YZ \cdot XZ} \geq \sqrt{\frac{2}{3}n \cdot (C \cdot dn)^2} = \sqrt{\frac{2}{3}}C \cdot dn^{3/2}$ , which proves the first lower bound.
- Assume that every vertex has aspect ratio at most  $r$  ( $r \geq 1$ ). In particular, therefore,  $Z(v) \leq rX(v)$  and  $Z(v) \leq rY(v)$  for every vertex  $v$ . Since the surface of  $v$  is at least  $\deg(v)$ , we have  $\deg(v) \leq 2(X(v)Y(v) + Y(v)Z(v) + X(v)Z(v)) \leq 6rX(v)Y(v)$ . Thus, every vertex  $v$  that intersects the  $(Z = Z_0)$  has  $X(v)Y(v) \geq \deg(v)/6r = d/6r$ . Since the  $(Z = Z_0)$ -plane intersects at least  $\frac{2}{3}n$  vertices, and these intersections are disjoint, there is at least  $\frac{2}{3}n \cdot d/6r$  grid points in the  $(Z = Z_0)$ -plane. Hence  $XY \geq \frac{1}{9} \cdot dn/r$ .



Therefore

$$XYZ = \sqrt{XY \cdot YZ \cdot XZ} \geq \sqrt{\frac{1}{9} \cdot dn/r \cdot (C \cdot dn)^2} = \frac{1}{3}C \cdot (dn)^{3/2}/\sqrt{r},$$

which proves the second lower bound.

- Assume now that the drawing is strictly  $\alpha$ -degree-restricted for some  $\alpha \geq 1$ . If  $X > Cn/\alpha$  and  $Y > Cn/\alpha$ , then  $XY > C^2n^2/\alpha^2$ , and  $XYZ = \sqrt{XY \cdot YZ \cdot XZ} \geq \sqrt{(C^2 \cdot n^2/\alpha^2) \cdot (C \cdot dn)^2} = C^2 \cdot dn^2/\alpha \geq C^2 \cdot (dn)^{3/2}/\alpha$  since  $d \leq n$ , which proves the claim.

So assume  $Y \leq Cn/\alpha$ ; the case  $X \leq Cn/\alpha$  is similar. Since the drawing is strictly  $\alpha$ -degree-restricted, the surface of every vertex  $v$  is at most  $\alpha \cdot \deg(v) = \alpha d$ , and therefore  $Z(v) < \alpha d/4$ .

Define  $Z_- = Z_0 - \alpha d/4$  and  $Z_+ = Z_0 + \alpha d/4$ . A point is *inside* if its  $Z$ -coordinate  $z$  satisfies  $Z_- < z < Z_+$ , and *outside* otherwise. Every vertex  $v$  in  $S$  and  $T$  crosses the  $(Z = Z_0)$ -plane. Since  $Z(v) < \alpha d/4$ ,  $v$  crosses neither the  $(Z = Z_-)$ -plane nor the  $(Z = Z_+)$ -plane, and thus all ports of  $v$  are inside.

Define  $X^* = X' + \frac{1}{2}$ ; we have shown above that at least  $C \cdot dn$  edges cross the  $(X = X^*)$ -plane. Of these, at least  $C \cdot dn - Y \cdot \alpha d/2$  edges cross the  $(X = X^*)$ -plane at an outside point, because there are at most  $Y \cdot \alpha d/2$  inside points with integer  $Y$ - and  $Z$ -coordinate on the  $(X = X^*)$ -plane.

Each of these  $C \cdot dn - Y \cdot \alpha d/2$  edges start at a vertex in  $S$  (and hence at an inside point), cross the  $(X = X^*)$ -plane at an outside point, and end at a vertex in  $T$  (also at an inside point). This implies that each edge crosses either the  $(Z = Z_-)$ -plane or the  $(Z = Z_+)$ -plane at least twice. Thus these two planes together have at least  $2(C \cdot dn - Y \cdot \alpha d/2)$  points with integral  $X$ - and  $Y$ -coordinate. Therefore  $XY \geq C \cdot dn - Y \cdot \alpha d/2 \geq C \cdot dn/2$  since  $Y \leq Cn/\alpha$ , and  $XYZ = \sqrt{XY \cdot YZ \cdot XZ} \geq \sqrt{\frac{1}{2}C \cdot dn \cdot (C \cdot dn)^2} = \frac{1}{\sqrt{2}}C^{3/2} \cdot (dn)^{3/2}$ .  $\square$

Using this theorem, we construct the lower bound for arbitrary values of  $m$  and  $n$ , as long as both are large enough. Before proving this, we need to show that primes congruent to 1 modulo 4 are frequent.

LEMMA 4. *There exist constants  $k \geq 2$  and  $x_1 \geq k$  such that for all  $x \geq x_1$ , the interval  $[(1/k)x, x]$  contains a prime number  $p$  with  $p \equiv 1 \pmod{4}$ .*

PROOF. Denote by  $\pi_{4,1}(x)$  the number of primes  $p \leq x$  that satisfy  $p \equiv 1 \pmod{4}$ . A famous theorem by de la Vallée Poussin establishes that

$$\pi_{4,1}(x) = \theta \left( \frac{1}{\varphi(4)} \cdot \frac{x}{\log x} \right),$$

where  $\varphi$  is Euler's function, in particular,  $\varphi(4) = 2$ . See, for example, [15] for a proof.

Let  $c_1, c_2, x_0$  be constants such that  $c_2 \geq c_1$  and

$$c_1 \cdot \frac{x}{\log x} \leq \pi_{4,1}(x) \leq c_2 \cdot \frac{x}{\log x}$$

for all  $x \geq x_0$ . Let  $k = 2c_2/c_1$ . Choose  $x_1 \geq k \cdot x_0$  so that  $\log(k) \leq \frac{1}{4} \log(x)$  and  $x/\log x \geq 3/c_1$  for all  $x \geq x_1$ . Then

$$\pi_{4,1}(x) - \pi_{4,1}\left(\frac{1}{k}x\right) \geq c_1 \frac{x}{\log x} - c_2 \frac{(c_1/2c_2)x}{\log(1/k) + \log x} \geq \frac{c_1 x}{\log x} - \frac{\frac{1}{2}c_1 x}{\frac{3}{4} \log x} \geq \frac{1}{3} \frac{c_1 x}{\log x} \geq 1.$$

Hence there is at least one prime number  $\equiv 1 \pmod{4}$  between  $(1/k)x$  and  $x$ .  $\square$

To establish this lower bound, we prove that for sufficiently large  $n$  and sufficiently large  $m$  there exists a graph  $G$  with  $n$  vertices and  $m$  edges that has an induced subgraph  $G'$  that satisfies the conditions of Theorem 1. Moreover,  $G'$  is asymptotically as big as  $G$ , i.e.,  $G'$  has  $\theta(n)$  vertices and  $\theta(m)$  edges.  $G'$  is also  $d'$ -regular, and vertices in  $G'$  have degree  $\theta(d')$  in  $G$ . The precise conditions are as follows:

LEMMA 5. *Let  $n \geq \max\{32k^4, 2x_1^2\}$  and  $m \geq \max\{\frac{1}{4}(144x_1 + 1)n, \frac{145}{4}kn\}$ , where  $k \geq 2$  and  $x_1 \geq k$  are the constants of Lemma 4. Then there exists a simple graph  $G$  with  $n$  vertices and  $m$  edges that has a  $d'$ -regular subgraph  $G'$  with  $n'$  vertices and  $m'$  edges such that:*

- $G'$  satisfies the conditions of Theorem 1.
- $n \geq n' \geq n/8k^2$ .
- $m \geq m' = d'n'/2 \geq m/64k^4$ .
- Every vertex of  $G'$  has degree at most  $8k^2 d'$  in  $G$ .

PROOF. The proof splits into two cases, depending on the size of  $m$ . If  $m$  is big enough, then  $G'$  can be a complete graph; if  $m$  is small, then we use a Ramanujan graph for  $G'$ . In both cases we “pad”  $G'$  with additional vertices and edges to achieve the desired number of vertices and edges.

*Case 1:  $m \geq n^2/64k^4$ .* In this case let  $G'$  be the complete graph on  $n' = \lceil n/8k^2 \rceil + 1$  vertices.

Clearly,  $G'$  as a complete graph satisfies the conditions of Theorem 1. Also,  $n/8k^2 + 1 \leq n' \leq n/8k^2 + 2 \leq n$ .  $G'$  has  $m' = \binom{n'}{2}$  edges, which is not too many, since  $m' \leq (n')^2/2 \leq (n/8k^2 + 2)^2/2 = n^2/128k^4 + n/4k^2 + 2 \leq n^2/64k^4 \leq m$  by  $n \geq 32k^4$ . Also,  $m' = \binom{n'}{2} \geq (n/8k^2)^2/2 = 1/64k^4 \cdot n^2/2 \geq 1/64k^4 \cdot m$  since  $m \leq \binom{n}{2}$ .

To obtain  $G$ , add  $n - n'$  vertices to  $G'$ , and add  $m - m'$  arbitrary edges such that the resulting graph is simple; this is possible since  $m \leq \binom{n}{2}$ . The maximum degree of  $G$  is at most  $n - 1$ .  $G'$  is  $d'$ -regular with  $d' = n' - 1 \geq n/8k^2$ . Hence the degree (in  $G$ ) of any vertex in  $G'$  is at most  $n - 1 \leq n \leq 8k^2 d'$ .

*Case 2:  $m < n^2/64k^4$ .* In this case,  $G'$  will be a Ramanujan graph  $G_{p,q}$  for some carefully chosen primes  $p$  and  $q$ .

Let  $q' = \frac{1}{2} + \sqrt{n/2 + \frac{1}{4}}$ , which implies that  $q'(q' - 1) = n/2$ ,  $q' \leq \sqrt{n/2} + 1$ , and  $q' \geq \sqrt{n/2} \geq x_1$ . Find a prime  $q$  with  $q \equiv 1 \pmod{4}$  such that  $(1/k)q' \leq q \leq q'$ ; this exists by Lemma 4. Note that  $q(q - 1) \leq q'(q' - 1) = n/2$ . Also, since  $n \geq 32k^4$  we have  $\sqrt{n/2} \leq n/8k^2$  and  $1 \leq n/8k^2$ . Hence  $\sqrt{n/2} + 1 \leq n/4k^2$  and  $q(q - 1) \geq q^2 - q \geq (q')^2/k^2 - q' \geq n/2k^2 - (\sqrt{n/2} + 1) \geq n/4k^2$ .

Let  $p' = 2m/q(q-1) - 1$ ; since  $m \geq \frac{1}{4}(144x_1 + 1)n$  and  $q(q-1) \leq n/2$  this implies that  $p' \geq 144x_1$ . Find a prime  $p$  with  $p \equiv 1 \pmod{4}$  such that  $(1/k)p' \leq p \leq p'$ ; this exists by Lemma 4. We have  $p \geq (1/k)p' \geq (1/k)144x_1 \geq 144$ . Also, since  $m \leq n^2/64k^4$  and  $q(q-1) \geq n/4k^2$  we have

$$p \leq p' < \frac{2n^2}{64k^4} \frac{1}{q(q-1)} \left( \frac{q(q-1)}{n/4k^2} \right)^2 = \frac{q(q-1)}{2}.$$

Let  $G'$  be the graph  $G_{p,q}$  for primes  $p$  and  $q$  as defined in Lemma 1, and suppose it has  $n'$  vertices and  $m'$  edges. By Lemma 1,  $G'$  satisfies the conditions of Theorem 1. Also,

$$n \geq \frac{n}{2} \geq q(q-1) \geq n' \geq \frac{1}{2}q(q-1) \geq \frac{n}{8k^2}.$$

By definition  $G'$  is  $d'$ -regular with  $d' = p + 1 \leq p' + 1$ , thus  $m'$  satisfies

$$m' = \frac{1}{2}(p+1)n' \leq \frac{1}{2}(p'+1)q(q-1) = m$$

and

$$m' = \frac{1}{2}(p+1)n' \geq \frac{1}{2k}(p'+k)\frac{1}{2}q(q-1) \geq \frac{1}{2k}m.$$

Create  $G$  by adding  $n - n'$  vertices to  $G'$ . Since  $m < n^2/64k^4$  we have  $2m \leq n^2/32 \leq (n/2 - 1)^2 \leq (n - q(q-1) - 1)^2 \leq (n - n' - 1)^2$ , hence  $m \leq \binom{n-n'}{2}$  and we can add  $m - m'$  edges between the  $n - n'$  added vertices in such a way that  $G$  is simple. Note that no incident edges were added to any vertex in  $G'$ ; in particular, the degree of vertices in  $G'$  remains  $d'$ . This completes the proof.  $\square$

Now we use these graphs to prove the lower bounds for almost any value of  $n$  and  $m$ .

**LEMMA 6.** *Let  $C \leq 1$  be the Ramanujan constant and let  $k \geq 2$  and  $x_1$  be the constants of Lemma 4. Then for any  $n \geq \max\{32k^4, 2x_1^2\}$  and  $m \geq \max\{\frac{1}{4}(144x_1 + 1)n, \frac{145}{4}kn\}$ , there exists a graph  $G$  with  $n$  vertices and  $m$  edges such that:*

- $\text{vol}(G, \infty, \infty) \geq \sqrt{2}/(384k^5) \cdot C^{3/2} \cdot m\sqrt{n}$ .
- $\text{vol}(G, r, \infty) \geq \sqrt{2}/(768k^6) \cdot C^{3/2} \cdot m^{3/2}/\sqrt{r}$ .
- $\text{vol}(G, \infty, \alpha) \geq \sqrt{2}/(6144k^8) \cdot C^2 \cdot m^{3/2}/\alpha$ .

**PROOF.** Let  $G$  be the graph of Lemma 5. Any drawing of  $G$  contains a drawing of  $G'$ , and thus has volume at least  $\frac{1}{3}C^{3/2}d'(n')^{3/2}$  by Theorem 1. Using the known bounds on  $n'$  and  $m'$  from Lemma 5, we obtain a lower bound on the volume of at least

$$\frac{1}{3}C^{3/2} \cdot 2m' \cdot \sqrt{n'} \geq \frac{2}{3}C^{3/2} \cdot m/64k^4 \cdot \sqrt{n/8k^2} = \sqrt{2}/(384k^5) \cdot C^{3/2} \cdot m\sqrt{n},$$

which proves the first claim.

If the drawing of  $G$  has aspect ratios at most  $r$ , then so does the drawing of  $G'$ , which thus has volume at least  $\frac{1}{3}C^{3/2}(d'n')^{3/2}/\sqrt{r}$ , which is at least

$$\frac{1}{3}C^{3/2} \cdot (2m')^{3/2}/\sqrt{r} \geq \sqrt{8}/3 \cdot C^{3/2} \cdot (m/64k^4)^{3/2}/\sqrt{r} = \sqrt{2}/(768k^6) \cdot C^{3/2} \cdot m^{3/2}/\sqrt{r};$$

this proves the second claim.

If the drawing of  $G$  is strictly  $\alpha$ -degree-restricted, then every vertex in  $G'$  has surface at most  $\alpha 8k^2 d'$ . Thus the drawing of  $G'$  is strictly  $\alpha 8k^2$ -degree-restricted, and has volume at least  $\frac{1}{3}C^2(d'n')^{3/2}/\alpha 8k^2$ , which is at least

$$1/24k^2 \cdot C^2 \cdot (2m')^{3/2}/\alpha \geq \sqrt{2}/12k^2 \cdot C^2 \cdot (m/64k^4)^{3/2}/\alpha = \sqrt{2}/(6144k^8) \cdot C^2 \cdot m^{3/2}/\alpha;$$

this proves the third claim.  $\square$

From Lemma 6 we can conclude the main result of this section.

**THEOREM 2.** *We have the following lower bounds:*

- $vol(n, m, \infty, \infty) = \Omega(m\sqrt{n})$ .
- $vol(n, m, r, \infty) = \Omega(m^{3/2}/\sqrt{r})$ .
- $vol(n, m, \infty, \alpha) = \Omega(m^{3/2}/\alpha)$ .

**3. Constructions.** In the following we give two constructions. The first creates degree-restricted cube-drawings with asymptotically optimal volume. The second creates drawings without restrictions on vertex boxes; again the drawings produced have asymptotically optimal volume.


**3.1. Cube-Drawings.** In the following algorithm for producing orthogonal drawings, each vertex  $v$  is initially represented by a square of size  $\mathcal{O}(\sqrt{\deg(v)}) \times \mathcal{O}(\sqrt{\deg(v)})$  in the  $(Z = 0)$ -plane. (The algorithm by Hagihara et al. [16] is similar in spirit, but uses squares of size  $\mathcal{O}(\sqrt{\Delta})$  for each vertex, and hence results in a drawing with  $\mathcal{O}((n\Delta)^{3/2})$  volume.) Edges are then routed either above or below the  $(Z = 0)$ -plane in a similar manner to the COMPACT point-drawing algorithm of Eades et al. [13]. Finally, the vertices are extended in the  $Z$ -dimension to form cubes.

**Algorithm** OPTIMAL VOLUME CUBE-DRAWING

*Input:* graph  $G = (V, E)$ .

*Output:* orthogonal drawing of  $G$ .

1. Initially represent each vertex  $v \in V$  by a square  $S_v$  with side length  $2 \lceil \sqrt{\lceil \deg(v)/2 \rceil + 1} \rceil$ .
2. Position the squares  $\{S_v : v \in V\}$  in the  $(Z = 0)$ -plane with the square-packing algorithm of Kleitman and Krieger [17]. Note that squares may touch, and since all squares have even side length, we may assume that all corners of the squares have even coordinates.
3. For each vertex  $v \in V$ , remove the top two rows from  $S_v$  and the two rightmost columns from  $S_v$ . Vertices are now disjoint; see Figure 1.
4. Pair the odd degree vertices in  $G$ , and add an edge between the paired vertices. All vertices now have even degrees. Orient the edges of  $G$  and alternately label the edges “+” and “−” by following an Eulerian tour of  $G$ . Remove the added edges.



**Fig. 1.** Positioning the vertices with a square-packing.

5. Assign each edge  $vw \in E$  labelled “+” unique  $Z^+$  ports at  $v$  and  $w$  both with an even  $X$ -coordinate and an even  $Y$ -coordinate.
6. Construct a graph  $H = (V_H, E_H)$  with vertex set  $V_H$  corresponding to the edges of  $G$  labelled “+”. For oriented edges  $vw, xy \in E$ , add the edge  $\{vw, xy\}$  to  $E_H$  if the port assigned to  $vw$  at  $v$  is in the same column as the port assigned to  $xy$  at  $x$ , or the port assigned to  $vw$  at  $w$  is in the same row as the port assigned to  $xy$  at  $y$ .
7. Determine a proper vertex-colouring of  $H$  using the sequential greedy algorithm (with colours  $\{1, 2, \dots, \Delta(H) + 1\}$ ). For each vertex of  $H$  coloured  $i$  corresponding to an edge  $vw$  in  $E$ , set the *height*  $h(vw) \leftarrow i$ .
8. For each oriented edge  $vw \in E$  labelled “+”, construct an edge route for  $vw$  as follows. Suppose the ports on  $v$  and  $w$  assigned to  $vw$  have coordinates  $(v_X, v_Y, 0)$  and  $(w_X, w_Y, 0)$ , respectively. Route the edge  $vw$  with one of the following 4- or 6-bend routes, as illustrated in Figure 2:
  - $v_X = w_X$ :  $(v_X, v_Y, 0) \rightarrow (v_X, v_Y, 2h(vw)) \rightarrow (v_X + 1, v_Y, 2h(vw)) \rightarrow (v_X + 1, w_Y, 2h(vw)) \rightarrow (v_X, w_Y, 2h(vw)) \rightarrow (v_X, w_Y, 0) = (w_X, w_Y, 0)$ .
  - $v_Y = w_Y$ :  $(v_X, v_Y, 0) \rightarrow (v_X, v_Y, 2h(vw) + 1) \rightarrow (v_X, v_Y + 1, 2h(vw) + 1) \rightarrow (w_X, v_Y + 1, 2h(vw) + 1) \rightarrow (w_X, v_Y, 2h(vw) + 1) \rightarrow (w_X, v_Y, 0) = (w_X, w_Y, 0)$ .
  - $v_X \neq w_X$  and  $v_Y \neq w_Y$ :  $(v_X, v_Y, 0) \rightarrow (v_X, v_Y, 2h(vw)) \rightarrow (v_X + 1, v_Y, 2h(vw)) \rightarrow (v_X + 1, w_Y + 1, 2h(vw)) \rightarrow (v_X + 1, w_Y + 1, 2h(vw) + 1) \rightarrow (w_X, w_Y + 1, 2h(vw) + 1) \rightarrow (w_X, w_Y, 2h(vw) + 1) \rightarrow (w_X, w_Y, 0)$ .
9. Repeat Steps 5–8 for the edges labelled “-” assigning  $Z^-$  ports and constructing edge routes below the  $(Z = 0)$ -plane in an analogous manner.
10. So that each vertex is enlarged into a cube, insert enough  $Z$ -planes at  $Z = 0$ , extend the side of each vertex parallel to the  $Z$ -axis, and possibly lengthen incident edges labelled “-”.




Fig. 2. Routing edges above the  $(Z = 0)$ -plane.

**THEOREM 3.** *Algorithm OPTIMAL VOLUME CUBE-DRAWING determines a 12-degree-restricted cube-drawing of any loopless graph  $G$  in  $\mathcal{O}(m^{3/2})$  time, with  $\mathcal{O}(m^{3/2})$  volume (assuming  $m \geq n$ ), and at most six bends per edge route.*

**PROOF.** After Step 3.1 the square  $S_v$  has side length  $2 \left\lceil \sqrt{\lceil \deg(v)/2 \rceil + 1} \right\rceil - 2$ . Since the corners of the square have even coordinates, the number of  $Z^+$  ports on  $S_v$  with even  $X$ - and  $Y$ -coordinates is at least

$$\left\lceil \sqrt{\lceil \deg(v)/2 \rceil + 1} \right\rceil^2 \geq \lceil \deg(v)/2 \rceil + 1.$$

At most  $\lceil \deg(v)/2 \rceil + 1$  edges incident to  $v$  are labelled “+”. (In fact, all vertices  $v$ , except the starting vertex in the Eulerian tour, have at most  $\lceil \deg(v)/2 \rceil$  incident edges labelled “+”.) Similarly, at most  $\lceil \deg(v)/2 \rceil + 1$  edges incident to  $v$  are labelled “−”. Thus there are enough ports on  $v$ .

If a unit-length edge segment intersects another edge route then so does one of the adjacent non-unit-length edge segments. Therefore, to show that the drawing is crossing-free, we need only show that non-unit-length edge segments do not intersect, and consider only such segments in the following. Vertical segments cannot intersect because ports are assigned to unique edges.  $X$ -segments have an odd  $Z$ -coordinate and  $Y$ -segments have an even  $Z$ -coordinate, thus an  $X$ -segment cannot intersect a  $Y$ -segment. A vertical segment has even  $X$ - and  $Y$ -coordinates, an  $X$ -segment has an odd  $Y$ -coordinate, and a  $Y$ -segment has an odd  $X$ -coordinate. Thus a vertical segment cannot intersect an  $X$ -segment or a  $Y$ -segment. Two  $Y$ -segments can only intersect if they overlap. Since edge routes originating in the same column have different heights, two  $Y$ -segments cannot intersect. Similarly, two  $X$ -segments can only intersect if originating in the same row, and in this case they have different heights, thus they cannot intersect. Hence no two edge routes can intersect.

The total area of the squares  $\{S_v : v \in V\}$  (before Step 3.1) is

$$\sum_v \left( 2 \left\lceil \sqrt{\lceil \deg(v)/2 \rceil + 1} \right\rceil \right)^2 \leq \sum_v \left( 2 \deg(v) + 4\sqrt{2(\deg(v) + 3)} + 10 \right),$$

which is

$$4m + \mathcal{O} \left( n + \sum_v \sqrt{\deg(v)} \right).$$

By the Cauchy–Schwarz inequality,  $\sum_v \sqrt{\deg(v)} \leq \sqrt{2nm}$ . Since we assume that  $m \geq n$  it follows that  $n \leq \sqrt{nm}$  and the total area of squares is therefore  $4m + \mathcal{O}(\sqrt{nm})$ .

The algorithm of Kleitman and Krieger [17] packs squares with a total area of 1 in a  $2/\sqrt{3} \times \sqrt{2}$  rectangle. Thus the squares  $\{S_v : v \in V\}$  can be packed in a rectangle with size

$$\left( 4\sqrt{\frac{m}{3}} + \mathcal{O}((nm)^{1/4}) \right) \times \left( 2\sqrt{2m} + \mathcal{O}((nm)^{1/4}) \right).$$

Hence the maximum degree of  $H$  is

$$\Delta(H) \leq \left( \frac{4}{\sqrt{3}} + 2\sqrt{2} \right) \sqrt{m} + \mathcal{O}((nm)^{1/4}).$$

A greedy vertex-colouring of  $H$  requires at most  $\Delta(H) + 1$  colours. Thus the height of the drawing above the  $(Z = 0)$ -plane, and the height below the vertices is at most

$$\left( \frac{8}{\sqrt{3}} + 4\sqrt{2} \right) \sqrt{m} + \mathcal{O}((nm)^{1/4}).$$

The height of the vertices is  $\max_v 2 \left\lceil \sqrt{(\deg(v) + 1)/2} \right\rceil \leq \sqrt{2\Delta(G)} + \mathcal{O}(1) \leq \sqrt{2m} + \mathcal{O}(1)$ ,

and thus the height of the drawing is at most

$$\left( \frac{16}{\sqrt{3}} + 9\sqrt{2} \right) \sqrt{m} + \mathcal{O}((nm)^{1/4}).$$

The bounding box is therefore at most

$$\left( 4\sqrt{\frac{m}{3}} + \mathcal{O}((nm)^{1/4}) \right) \times \left( 2\sqrt{2m} + \mathcal{O}((nm)^{1/4}) \right) \times \left( \left( \frac{16}{\sqrt{3}} + 9\sqrt{2} \right) \sqrt{m} + \mathcal{O}((nm)^{1/4}) \right).$$

A simple calculation establishes that the bounding box volume is at most

$$\left( \frac{144}{\sqrt{3}} + \frac{128\sqrt{2}}{3} \right) m^{3/2} + \mathcal{O}(m(nm)^{1/4}) \leq 144m^{3/2} + \mathcal{O}(m(nm)^{1/4}) = \mathcal{O}(m^{3/2}).$$

The time-consuming stage of the algorithm is the vertex colouring of  $H$ . This can be computed in  $\mathcal{O}(|E_H|) = \mathcal{O}(|V_H| \Delta(H)) = \mathcal{O}(m\sqrt{m}) = \mathcal{O}(m^{3/2})$  time. The surface of a vertex  $v$  is

$$6 \left( 2 \left\lceil \sqrt{(\deg(v) + 1)/2} \right\rceil \right)^2 \leq 12 \cdot \deg(v) + \mathcal{O}(\sqrt{\deg(v)}).$$

Thus each vertex is 12-degree-restricted. By construction, there are at most six bends per edge route.  $\square$

Biedl and Chan [6] have developed a technique based on edge-colouring a certain bipartite graph to implement Steps 3.1 and 3.1 of Algorithm OPTIMAL VOLUME CUBE-DRAWING more efficiently. With this technique, the time complexity reduces to  $\mathcal{O}(m \log m)$ , and the volume of the produced drawings decreases to  $\approx 83m^{3/2} + \mathcal{O}(m(nm)^{1/4})$ .

The number of bends per edge can be reduced to five if we allow for an increase in the volume.

**THEOREM 4.** *Every loopless graph has a 12-degree-restricted cube-drawing, which can be computed in  $\mathcal{O}(m)$  time, with  $\mathcal{O}(m^2)$  volume, and at most five bends per edge route.*

**PROOF.** The claimed drawing is obtained by modifying Algorithm OPTIMAL VOLUME CUBE-DRAWING in two ways. Number the edges  $e_1, \dots, e_m$ , and set  $h(e_i) = i/2$  for all  $i$  (and hence omit Steps 6 and 7). Then in Step 8, in the case that  $v_X \neq w_X$  and  $v_Y \neq w_Y$ , omit the middle segment, and apply the routing  $(v_X, v_Y, 0) \rightarrow (v_X, v_Y, 2h(vw)) \rightarrow (v_X + 1, v_Y, 2h(vw)) \rightarrow (w_X, w_Y + 1, 2h(vw)) \rightarrow (w_X, w_Y, 2h(vw)) \rightarrow (w_X, w_Y, 0)$ . Since  $h(e_i) \neq h(e_j)$  for all  $i \neq j$ , one can easily verify that this gives a crossing-free drawing with bounding box  $\mathcal{O}(\sqrt{m}) \times \mathcal{O}(\sqrt{m}) \times \mathcal{O}(m)$ .  $\square$

**3.2. Drawings with an Unbounded Aspect Ratio.** We now show how to create drawings of a simple graph that have volume  $\mathcal{O}(m\sqrt{n})$ , which is optimal. The vertices have an unbounded aspect ratio and are not necessarily degree-restricted. can easily

The following algorithm initially represents vertices by points or line segments in the ( $Z = 0$ )-plane, and edges are routed above this plane. The vertices are then extended in the  $Z$ -dimension to form lines or rectangles.

**Algorithm OPTIMAL VOLUME BOX-DRAWING**


*Input:* simple graph  $G = (V, E)$ .

*Output:* orthogonal drawing of  $G$ .

1. Let  $N = \lceil \sqrt{n} \rceil$ , and define  $V_{\text{big}} = \{v \in V : \deg(v) > 4m/N\}$  and  $V_{\text{small}} = V \setminus V_{\text{big}}$ .  
Define  $E_{\text{big}}$  to be the set of edges with both endpoints in  $V_{\text{big}}$ ,  $E_{\text{cut}}$  to be the set of edges with exactly one endpoint in  $V_{\text{big}}$ , and  $E_{\text{small}}$  to be the set of edges with both endpoints in  $V_{\text{small}}$ .
2. Let  $V_{\text{big}} = \{w_1, \dots, w_l\}$ . For each vertex  $w_j \in V_{\text{big}}$ , define  $k(w_j) = \lceil \deg(w_j)N/4m \rceil$ , and initially represent  $w_j$  by the line segment with endpoints at

$$\left( 2 \sum_{i=1}^{j-1} k(w_i), -2, 0 \right) \text{ and } \left( 2 \left( \sum_{i=1}^j k(w_i) - 1 \right), -2, 0 \right).$$





**Fig. 3.** Placing points and segments of vertices. Vertices  $w_1, w_2, w_3$  belong to  $V_{\text{big}}$ .

Note that  $w_j$  has  $k(w_j) \geq 2$  grid points with an even  $X$ -coordinate. When  $j = 1$  the sum on the left is taken to be 0.

3. Add extra vertices of degree 0 to  $V_{\text{small}}$  so that  $V_{\text{small}}$  has exactly  $N^2$  vertices (simply to ease the description). Let  $V_{\text{small}} = (v_0, v_1, v_2, \dots, v_{N^2-1})$  sorted by non-increasing degree. For each original vertex  $v_k \in V_{\text{small}}, 0 \leq k \leq N^2 - 1$ , initially represent  $v_k$  by the point  $(2(i + j) \bmod 2N, 2j, 0)$ , where  $k = iN + j$  for the unique pair  $i, j$  with  $0 \leq i, j \leq N - 1$ . See Figure 3.
4. Orient each edge in  $E_{\text{cut}}$  from the endpoint in  $V_{\text{big}}$  to the other endpoint. Orient all edges in  $E_{\text{small}}$  arbitrarily.
5. For each oriented edge  $e = vw \in E_{\text{cut}}$ , assign to  $e$  a  $Z^+$ -port at  $v$  with even coordinates, such that at most  $\lceil 4m/N \rceil$  edges are assigned to any  $Z^+$ -port of any vertex in  $V_{\text{big}}$ . Assign the unique  $Z^+$ -port at  $w$  to  $e$ . For each edge  $e = vw \in E_{\text{small}}$ , assign to  $e$  the unique  $Z^+$ -ports at  $v$  and  $w$ .
6. Construct a graph  $H = (V_H, E_H)$  with vertex set  $V_H$  corresponding to the edges of  $E_{\text{cut}} \cup E_{\text{small}}$ . For oriented edges  $vw, xy \in E_{\text{cut}} \cup E_{\text{small}}$ , add the edge  $\{vw, xy\}$  to  $E_H$  if the port assigned to  $vw$  at  $v$  is in the same column as the port assigned to  $xy$  at  $x$ , or the port assigned to  $vw$  at  $w$  is in the same row as the port assigned to  $xy$  at  $y$ .
7. Determine a proper vertex-colouring of  $H$  using the sequential greedy algorithm (with colours  $\{1, 2, \dots, \Delta(H) + 1\}$ ). For each vertex of  $H$  coloured  $i$  corresponding to an edge  $vw \in E_{\text{cut}} \cup E_{\text{small}}$ , set the *height*  $h(vw) \leftarrow i$ .
8. For each oriented edge  $vw \in E_{\text{cut}} \cup E_{\text{small}}$  construct an edge route for  $vw$  exactly as described in Step 3.1 of Algorithm OPTIMAL VOLUME CUBE-DRAWING.

9. Construct edge routes for edges in  $E_{\text{big}}$  by copying the 2-bend layout of the complete graph developed in [8]. More precisely, recall that  $|V_{\text{big}}| = l$ . It is possible to partition the edges of the complete graph  $K_l$ , and therefore  $E_{\text{big}}$ , into  $l$  matchings  $M_1, M_2, \dots, M_l$  such that if the edges  $(w_i, w_j)$ ,  $i < j$ , and  $(w_a, w_b)$ ,  $a < b$ , are in the same matching, and (say)  $i \leq a$ , then  $i < a < b < j$  or  $i < j < a < b$ . If the edge  $(w_i, w_j)$  with  $i < j$  belongs to matching  $M_k$ , and if the leftmost points of  $w_i$  and  $w_j$  have coordinates  $(x_i, -2, 0)$  and  $(x_j, -2, 0)$ , respectively, then route  $(w_i, w_j)$  as

$$\begin{aligned} (x_i, -2, 0) &\rightarrow (x_i, -2, k) \rightarrow (x_i, -2 - \lceil (j-i)/2 \rceil, k) \\ &\rightarrow (x_j, -2 - \lceil (j-i)/2 \rceil, k) \rightarrow (x_j, -2, k) \rightarrow (x_j, -2, 0). \end{aligned}$$

10. Enlarge points/lines representing vertices into lines/rectangles by extending their sides parallel to the  $Z$ -axis from the minimum to the maximum  $Z$ -coordinate of the drawing. For each edge  $vw \in E$ , clip the segment of  $vw$  incident to  $v$  if it is contained in the box representing  $v$  (and similarly at  $w$ ); see Figure 4.

To analyse the size of the resulting drawing, we need a bound on the maximum degree of  $H$ .

LEMMA 7. *The maximum degree of  $H$  in Algorithm OPTIMAL VOLUME BOX-DRAWING is at most  $16m/N - 1$ .*

PROOF. Let  $xy$  be an arbitrary oriented edge in  $E_{\text{cut}} \cup E_{\text{small}}$ . Any neighbour of  $xy$  in  $H$  must have a port in the same column as  $x$  or in the same row as  $y$ . Thus we want to study the number of edges that have a port in a specific row/column.




Fig. 4. A selection of edge routes in Algorithm OPTIMAL VOLUME BOX-DRAWING;  $e_1$ ,  $e_2$  and  $e_3$  belong to  $E_{\text{big}}$ ,  $e_4$  and  $e_5$  belong to  $E_{\text{cut}}$ , and  $e_6$  and  $e_7$  belong to  $E_{\text{small}}$ .

Assume that  $xy \in E_{\text{cut}}$ . By the direction of edges, we know that  $x \in V_{\text{big}}$  and  $y \in V_{\text{small}}$ . If  $vw$  is a neighbour of  $xy$  in  $H$ , because  $w$  and  $y$  have ports in the same row  $r$ , then row  $r$  contains vertices in  $V_{\text{small}}$ , and therefore this is *not* the row with  $Y$ -coordinate  $-2$ . If  $xy \in E_{\text{small}}$ , then both ports must be in a row that does not have  $Y$ -coordinate  $-2$  because both endpoints are in  $V_{\text{small}}$ . Hence no edge in  $H$  is added because of the row with  $Y$ -coordinate  $-2$ , which we can therefore ignore for considerations of the maximum degree of  $H$ .

Also, notice that by the way we assign ports to edges, there are at most  $\lceil 4m/N \rceil$  edges in  $E_{\text{cut}}$  assigned to the same  $Z^+$ -port of a vertex in  $V_{\text{big}}$  (i.e., to a  $Z^+$ -port with  $Y$ -coordinate  $-2$ ). To obtain a bound on the maximum degree, we thus need a bound on the number of edges in  $H$  resulting from a  $Z^+$ -port with a non-negative  $Y$ -coordinate. For future reference, let a *proper row/column* be the intersection of a row/column with the first quadrant, i.e., with the range of non-negative  $X$ - and  $Y$ -coordinates.

CLAIM. *For any proper row and any proper column, there are at most  $6m/N$  edges that use a  $Z^+$ -port in that row/column.*

We prove this by showing that the sum of the degrees of vertices placed in any one proper row/column is at most  $6m/N$ . Define for  $i = 0, \dots, N-1$  the  $i$ th *diagonal* to be the grid-points  $\{(2(i+k) \bmod 2N, 2k) : k = 0, \dots, N-1\}$ . In Step 3.2, vertices in  $V_{\text{small}}$  are positioned, in order of non-increasing degree, in the 0th diagonal from bottom to top, then in the 1th diagonal from bottom to top, and so on. This implies the following properties for each  $i$ ,  $0 \leq i \leq N-1$ :

- The vertex with the largest degree in the  $i$ th diagonal is at  $(2i, 0)$ , while the vertex with the smallest degree in this diagonal is at  $(2(i+N-1) \bmod 2N, 2(N-1))$ .
- The last vertex in the  $i$ th diagonal (i.e., the vertex at  $(2(i+N-1) \bmod 2N, 2(N-1))$ ) has degree no larger than the first vertex in the  $(i+1)$ th diagonal (i.e., the vertex at  $(2(i+1), 0)$ ).

For any proper row/column, define the *degree-sum* as the sum of the degrees of the vertices placed in this row/column. Denote the degree-sum of the row with  $Y$ -coordinate  $2i$  by  $R(i)$ .

Notice that each proper column and each proper row contains exactly one vertex from each diagonal, and no other vertices. Thus the degree-sum of each proper row/column can be at most the sum of the maximum degrees in each diagonal, and is at least the sum of the minimum degrees in each diagonal. By the first observation above, we know that the degree-sum of each proper row/column is at most  $R(0)$  and at least  $R(N-1)$ .

Also, by the second observation above, we know that  $R(0) \leq R(N-1) + \Delta_{\text{small}}$ , where  $\Delta_{\text{small}}$  is the maximum degree among all vertices in  $V_{\text{small}}$ , in particular,  $\Delta_{\text{small}} \leq 4m/N$  by the definition of  $V_{\text{small}}$ . This follows because each entry in  $R(0)$  is the first entry in some diagonal, and can be upper bounded by the last entry in the previous diagonal. The only exception is the first entry in the 0th diagonal, which has degree  $\Delta_{\text{small}}$ .

Now we can estimate

$$2m \geq \sum_{v \in V_{\text{small}}} \deg(v) = \sum_{i=0}^{N-1} R(i) \geq N \cdot R(N-1) \geq N \cdot R(0) - N \Delta_{\text{small}},$$

and therefore  $R(0) \leq 2m/N + \Delta_{\text{small}} \leq 6m/N$ . Since the degree-sum of each proper row/column is at most  $R(0)$ , the claim follows.

So any vertex in  $H$  (which corresponds to an edge in  $E_{\text{cut}} \cup E_{\text{small}}$ ) has at most  $6m/N - 1$  neighbours that use a  $Z^+$ -port in the same row, and at most  $6m/N - 1 + \lceil 4m/N \rceil$  neighbours that use a  $Z^+$ -port in the same column. Hence the maximum degree of  $H$  is at most  $16m/N - 1$ .  $\square$

**THEOREM 5.** *Algorithm OPTIMAL VOLUME BOX-DRAWING determines a drawing of any simple graph  $G$  in  $\mathcal{O}(m^2/\sqrt{n})$  time, with  $\mathcal{O}(m\sqrt{n})$  volume, and at most four bends per edge route.*

**PROOF.** First, we show that no edges overlap or intersect. Exactly as in Theorem 3, one shows that if two edges in  $E_{\text{cut}} \cup E_{\text{small}}$  have different ports at both endpoints, then the edges neither overlap nor cross. The same holds for any two edges in  $E_{\text{big}}$  as discussed in [8]. No edge of  $E_{\text{big}}$  can overlap or intersect an edge in  $E_{\text{cut}} \cup E_{\text{small}}$ , because they are separated by the  $(Y = 0)$ -plane.

If two edges  $xy$  and  $vw$  have a common port at one endpoint, say at  $x = v$ , then the edges do overlap, but only at the segment incident to  $x = v$ , which is parallel to the  $Z$ -axis. This segment will be clipped when extending the vertices in Step 3.2; hence there is no overlap in the final drawing.

Since  $V_{\text{small}}$  contains at most  $n \leq N^2$  vertices, the vertices in  $V_{\text{small}}$  can be placed in a  $2N \times 2N$  rectangle. Since  $\deg(v) > 4m/N$  for all vertices  $v \in V_{\text{big}}$ , we have

$$|V_{\text{big}}| \leq \sum_{v \in V_{\text{big}}} \frac{\deg(v)N}{4m} \leq \frac{N}{2}$$

and

$$\sum_{v \in V_{\text{big}}} k(v) = \sum_{v \in V_{\text{big}}} \left\lceil \frac{\deg(v)N}{4m} \right\rceil \leq \sum_{v \in V_{\text{big}}} \frac{\deg(v)N}{4m} + |V_{\text{big}}| \leq N,$$

thus the maximum  $X$ -coordinate of a vertex in  $V_{\text{big}}$  is  $2N$ , and we need at most  $2N$   $X$ -planes.

Since  $|V_{\text{big}}| \leq N/2$ , the edges in  $E_{\text{big}}$  have  $Y$ -coordinate  $\geq -2 - \lceil (N/2 - 1)/2 \rceil$ . Hence we need at most  $\frac{9}{4}N + 3$   $Y$ -planes.

In Step 5 a vertex  $v \in V_{\text{big}}$  has at most  $\deg(v)$  incident edges that need a  $Z^+$ -port in  $v$ , and there are  $k(v)$   $Z^+$ -ports at  $v$  with even coordinates. Thus, we can assign edges to ports such that each port has at most  $\lceil \deg(v)/k(v) \rceil \leq \lceil 4m/N \rceil$  edges assigned to it. Hence there are enough ports for the edges.

A greedy vertex colouring of  $H$  requires at most  $\Delta(H) + 1 \leq 16m/N$  colours, thus to route the edges in  $E_{\text{cut}} \cup E_{\text{small}}$ , we need at most  $2 \cdot 16m/N$   $Z$ -planes. To route the edges in  $E_{\text{big}}$ , we need at most  $|V_{\text{big}}| \leq N/2$   $Z$ -planes. Since  $m \geq n$ , we have  $32m/N \geq N/2$ , and the height of the drawing above the  $Z = 0$  plane is at most  $32m/N$ . The bounding box therefore has volume at most  $2N \times (\frac{9}{4}N + 3) \times 32m/N = 144mN + \mathcal{O}(m) = 144m\sqrt{n} + \mathcal{O}(m) = \mathcal{O}(m\sqrt{n})$ .

The time-consuming stage of the algorithm is the vertex colouring of  $H$ . This can be computed in  $\mathcal{O}(|E_H|) = \mathcal{O}(|V_H|\Delta(H)) = \mathcal{O}(m \cdot m/N) = \mathcal{O}(m^2/\sqrt{n})$  time.

Originally, there are at most six bends per edge route, and at most four bends per edge route for edges in  $V_{\text{big}}$ . During the clipping step, the first and last segment of each edge gets clipped. Hence every edge has at most four bends.  $\square$

The technique of Biedl and Chan [6] works for Steps 3.2 and 3.2 of Algorithm OPTIMAL VOLUME BOX-DRAWING as well, reducing the time complexity to  $\mathcal{O}(m \log n)$ , and decreasing the volume of the produced drawings to  $\approx 90m\sqrt{n} + \mathcal{O}(m)$ .

Note that we required  $G$  to be simple; this is necessary for the routing of the edges in  $E_{\text{big}}$ . It is not hard to show that if each edge in  $E_{\text{big}}$  has multiplicity  $\leq k$ , then all edges in  $E_{\text{big}}$  can be routed with  $\leq kN$   $Z$ -planes. Thus, as long as  $m = \Omega(kn)$ , the drawing still has  $\mathcal{O}(m\sqrt{n})$  volume.

In an analogous way to Theorem 4, the number of bends per edge can be reduced to three if we allow for an increase in the volume.

**THEOREM 6.** *Every loopless graph has a drawing, which can be computed in  $\mathcal{O}(m)$  time, with  $\mathcal{O}(mn)$  volume, and three bends per edge route.*

**PROOF.** The construction is a greatly simplified version of Algorithm OPTIMAL VOLUME BOX-DRAWING. In particular, there is now no need for  $V_{\text{big}}$  and no need to sort the vertices by degree. Number the edges  $e_1, \dots, e_m$ , and set  $h(e_i) = i/2$  for all  $i$ . Arbitrarily place the vertices at the points  $\{(2i, 2j, 0) : 1 \leq i, j \leq \lceil \sqrt{n} \rceil\}$ . Route each edge as before, except in the case that  $v_X \neq w_Y$  and  $v_Y \neq w_Y$ . Here the middle segment is omitted, and the edge  $vw$  is routed  $(v_X, v_Y, 2h(vw)) \rightarrow (v_X + 1, v_Y, 2h(vw)) \rightarrow (v_X + 1, w_Y + 1, 2h(vw)) \rightarrow (w_X + 1, w_Y + 1, 2h(vw)) \rightarrow (w_X, w_Y, 2h(vw))$ . Finally extend the points representing the vertices into  $Z$ -lines of length  $m$ . Since each edge lies in a unique  $Z$ -plane, one can easily verify that we obtain a crossing-free drawing with bounding box  $2 \lceil \sqrt{n} \rceil \times 2 \lceil \sqrt{n} \rceil \times m$ .  $\square$

Theorem 6 is particularly appropriate for multilayer VLSI as there are no vertical edge segments or “cross-cuts”; see [2].

**4. Conclusions and Open Problems.** In this paper we provided asymptotically matching upper and lower bounds on the volume of three-dimensional orthogonal box-drawings, under various restrictions on the shape of vertex boxes.

In particular, we showed that any algorithm to create three-dimensional orthogonal drawings that have bounded aspect ratios or are degree-restricted cannot do better than  $\Omega(m^{3/2})$  volume. Then we gave an algorithm that matches this bound, i.e., constructs three-dimensional degree-restricted orthogonal cube-drawings with  $\mathcal{O}(m^{3/2})$  volume.

If there are no restrictions on the drawing, then we showed that no algorithm can do better than  $\Omega(m\sqrt{n})$  volume. We gave a second algorithm that matches this bound, i.e., constructs three-dimensional orthogonal drawings with  $\mathcal{O}(m\sqrt{n})$  volume.

Thus, no more asymptotic improvements are possible for the volume of drawings. We do see room for improvement with respect to the number of bends per edge. In particular, (a) does every graph have a 5-bend degree-restricted cube-drawing with

$\mathcal{O}(m^{3/2})$  volume, and (b) does every graph have a 3-bend drawing with  $\mathcal{O}(m\sqrt{n})$  volume? Note that  $K_n$  does have a  $\mathcal{O}(n^{5/2}) = \mathcal{O}(m\sqrt{n})$  volume 3-bend drawing [8].

Table 1 suggests a trade-off between the number of bends per edge and the bounding box volume. Can such a trade-off be proved? What are lower bounds for drawings where edges are allowed to have at most  $k$  bends per edge, for  $k = 1, \dots, 5$ ? (Note that some graphs do not have a drawing without bends [8], [14], and lower bounds for drawings with one bend per edge were given in [4].)

## References

- [1] Mokhtar A. Aboelaze and Benjamin W. Wah. Complexities of layouts in three-dimensional VLSI circuits. *Inform. Sci.*, 55(1–3):167–188, 1991.
- [2] Alok Aggarwal, Maria Klawe, and Peter Shor. Multilayer grid embeddings for VLSI. *Algorithmica*, 6(1):129–151, 1991.
- [3] Noga Alon and Joel H. Spencer. *The Probabilistic Method*. Wiley, New York, 1992.
- [4] Therese Biedl. 1-Bend 3-D orthogonal box-drawings: two open problems solved. *J. Graph Algorithms Appl.*, 5(3):1–15, 2001.
- [5] Therese C. Biedl. Three approaches to 3D-orthogonal box-drawings. In Sue Whitesides, editor, *Proc. 6th International Symp. on Graph Drawing (GD '98)*, volume 1547 of Lecture Notes in Computer Science, pages 30–43. Springer, Berlin, 1998.
- [6] Therese Biedl and Timothy Chan. A note on 3D orthogonal graph drawing. *Discrete Appl. Math.*, to appear.
- [7] Therese C. Biedl and Michael Kaufmann. Area-efficient static and incremental graph drawings. In Rainer E. Burkhard and Gerhard J. Woeginger, editors, *Proc. 5th Annual European Symp. on Algorithms (ESA '97)*, volume 1284 of Lecture Notes in Comput. Science, pages 37–52. Springer, Berlin, 1997.
- [8] Therese C. Biedl, Thomas C. Shermer, Sue Whitesides, and Stephen K. Wismath. Bounds for orthogonal 3-D graph drawing. *J. Graph Algorithms Appl.*, 3(4):63–79, 1999.
- [9] M. Closson, S. Gartshore, John R. Johansen, and Stephen K. Wismath. Fully dynamic 3-dimensional orthogonal graph drawing. *J. Graph Algorithms Appl.*, 5(2):1–34, 2001.
- [10] Giuseppe Di Battista, Maurizio Patrignani, and Francesco Vargiu. A split&push approach to 3D orthogonal drawing. *J. Graph Algorithms Appl.*, 4(3):105–133, 2000.
- [11] Guisepe Di Battista, Peter Eades, Roberto Tamassia, and Ioannis G. Tollis. *Graph Drawing: Algorithms for the Visualization of Graphs*. Prentice-Hall, Englewood Cliffs, NJ, 1999.
- [12] Peter Eades, Charles Stirk, and Sue Whitesides. The techniques of Kolmogorov and Barzdin for three dimensional orthogonal graph drawings. *Inform. Process Lett.*, 60(2):97–103, 1996.
- [13] Peter Eades, Antonios Symvonis, and Sue Whitesides. Three dimensional orthogonal graph drawing algorithms. *Discrete Appl. Math.*, 103:55–87, 2000.
- [14] Sándor Fekete and Henk Meijer. Rectangle and box visibility graphs in 3D. *Internat. J. Comput. Geom. Appl.*, 9(1):1–27, 1999.
- [15] Ernests K. Fogels. An elementary proof of formulae of de la Vallée Poussin (in Russian). *Latvijas PSR Zinātņu Akad. Vestis*, 11(40):123–130, 1950.
- [16] Ken'ichi Hagihara, Nobuki Tokura, and Naoya Suzuki. Graph embedding on a three-dimensional model. *Systems Comput. Controls*, 14(6):58–66, 1983.
- [17] Daniel J. Kleitman and Michael M. Krieger. An optimal bound for two dimensional bin packing. In *Proc. 16th Annual Symp. on Foundations of Computer Science (FOCS '75)*, pages 163–168. IEEE, New York, 1975.
- [18] F. Thomson Leighton and Arnold L. Rosenberg. Three-dimensional circuit layouts. *SIAM J. Comput.*, 15(3):793–813, 1986.
- [19] Alexander Lubotzky, Ralph Phillips, and Peter Sarnak. Ramanujan graphs. *Combinatorica*, 8:261–277, 1988.
- [20] Gregory A. Margulis. Explicit group-theoretical constructions of combinatorial schemes and their applications to the design of expanders and superconcentrators. *Probl. Inf. Transm.*, 24(1):39–64, 1988.

- [21] Achilleas Papakostas and Ioannis G. Tollis. Algorithms for incremental orthogonal graph drawing in three dimensions. *J. Graph Algorithms Appl.*, 3(4):81–115, 1999.
- [22] Franco P. Preparata. Optimal three-dimensional VLSI layouts. *Math. Systems Theory*, 16:1–8, 1983.
- [23] Arnold L. Rosenberg. Three-dimensional VLSI: a case study. *J. Assoc. Comput. Mach.*, 30(2):397–416, 1983.
- [24] Avi Wigderson and David Zuckerman. Expanders that beat the eigenvalue bound: explicit construction and applications. In *Proc. 25th Annual ACM Symp. on Theory of Computing (STOC '93)*, pages 245–251, 1993.
- [25] David R. Wood. Multi-dimensional orthogonal graph drawing with small boxes. In Jan Kratochvil, editor, *Proc. 7th International Symp. on Graph Drawing (GD '99)*, volume 1731 of Lecture Notes in Computer Science, pages 311–222. Springer, Berlin, 1999.
- [26] David R. Wood. A new algorithm and open problems in three-dimensional orthogonal graph drawing. In Rajeev Raman and Jamie Simpson, editors, *Proc. 8th Australasian Workshop on Combinatorial Algorithms (AWOCA '97)*, pages 157–167. Curtin University of Technology, Australia, 1999.
- [27] David R. Wood. Bounded degree book embeddings and three-dimensional orthogonal graph drawing. In P. Mutzel, M. Jünger, and S. Leipert, editors, *Proc. 9th International Symp. on Graph Drawing (GD '01)*, volume 2265 of Lecture Notes in Computer Science, pages 312–327. Springer, Berlin, 2002.
- [28] David R. Wood. Optimal three-dimensional orthogonal graph drawing in the general position model. *Theoret. Comput. Sci.*, 299(1–3):151–178, 2003.
- [29] David R. Wood. Minimising the number of bends and volume in three-dimensional orthogonal graph drawings with a diagonal vertex layout. *Algorithmica*, 39(3):235–253, 2004.