

Planar decompositions and the crossing number of graphs with an excluded minor

David R. Wood and Jan Arne Telle

ABSTRACT. Tree decompositions of graphs are of fundamental importance in structural and algorithmic graph theory. Planar decompositions generalise tree decompositions by allowing an arbitrary planar graph to index the decomposition. We prove that every graph that excludes a fixed graph as a minor has a planar decomposition with bounded width and a linear number of bags.

The crossing number of a graph is the minimum number of crossings in a drawing of the graph in the plane. We prove that planar decompositions are intimately related to the crossing number. In particular, a graph with bounded degree has linear crossing number if and only if it has a planar decomposition with bounded width and linear order. It follows from the above result about planar decompositions that every graph with bounded degree and an excluded minor has linear crossing number.

Analogous results are proved for the convex and rectilinear crossing numbers. In particular, every graph with bounded degree and bounded tree-width has linear convex crossing number, and every $K_{3,3}$ -minor-free graph with bounded degree has linear rectilinear crossing number.

CONTENTS

1. Introduction	118
1.1. Complementary results	119
2. Graph decompositions	120
3. Manipulating decompositions	123
4. Planar decompositions and the crossing number	126
5. K_5 -minor-free graphs	130
6. Graphs embedded on a surface	134
7. H -minor-free graphs	137

Received April 21, 2006, Revised: May 22, 2007.

Mathematics Subject Classification. 05C62 (graph representations), 05C10 (topological graph theory), 05C83 (graph minors).

Key words and phrases. Graph drawing, crossing number, rectilinear crossing number, convex crossing number, outerplanar crossing number, graph decomposition, planar decomposition, tree decomposition, tree-width, tree-partition, tree-partition-width, planar partition, graph minor.

The research of David Wood is supported by a Marie Curie Fellowship of the European Community under contract 023865, and by the projects MEC MTM2006-01267 and DURSI 2005SGR00692.

8. Graph partitions	139
9. $K_{3,3}$ -minor-free graphs	141
Acknowledgements	142
References	142

1. Introduction

The *crossing number* of a graph¹ G , denoted by $\text{cr}(G)$, is the minimum number of crossings in a drawing² of G in the plane; see [36, 62, 91] for surveys. The crossing number is an important measure of the nonplanarity of a graph [87], with applications in discrete and computational geometry [60, 86] and VLSI circuit design [8, 49, 50]. In information visualisation, one of the most important measures of the quality of a graph drawing is the number of crossings [66, 65, 67].

Computing the crossing number is \mathcal{NP} -hard [39], and remains so for simple cubic graphs [45, 64]. Moreover, the exact or even asymptotic crossing number is not known for specific graph families, such as complete graphs [71], complete bipartite graphs [54, 69, 71], and cartesian products [1, 15, 41, 70].

Given that the crossing number seems so difficult, it is natural to focus on asymptotic bounds rather than exact values. The ‘crossing lemma’, conjectured by Erdős and Guy [36] and first proved³ by Leighton [49] and Ajtai et al. [5], gives such a lower bound. It states that

$$\text{cr}(G) \geq \|G\|^3/64|G|^2$$

for every graph G with $\|G\| \geq 4|G|$. Other general lower bound techniques that arose out of the work of Leighton [49, 50] include the bisection/cutwidth method [32, 59, 84, 85] and the embedding method [82, 84].

Upper bounds on the crossing number of general families of graphs have been less studied, and are the focus of this paper. Obviously $\text{cr}(G) \leq \binom{\|G\|}{2}$ for every graph G . A family of graphs has *linear*⁴ crossing number if

$$\text{cr}(G) \leq c|G|$$

¹We consider graphs G that are undirected, simple, and finite. Let $V(G)$ and $E(G)$ respectively be the vertex and edge sets of G . Let $|G| := |V(G)|$ and $\|G\| := |E(G)|$. For each vertex v of a graph G , let $N_G(v) := \{w \in V(G) : vw \in E(G)\}$ be the neighbourhood of v in G . The *degree* of v is $|N_G(v)|$. Let $\Delta(G)$ be the maximum degree of a vertex of G .

²A *drawing* of a graph represents each vertex by a distinct point in the plane, and represents each edge by a simple closed curve between its endpoints, such that the only vertices an edge intersects are its own endpoints, and no three edges intersect at a common point (except at a common endpoint). A *crossing* is a point of intersection between two edges (other than a common endpoint). A drawing with no crossings is *plane*. A graph is *planar* if it has a plane drawing.

³A remarkably simple probabilistic proof of the crossing lemma was found by Chazelle, Sharir and Welzl [4]. See [53, 58] for recent improvements.

⁴If the crossing number of a graph is linear in the number of edges then it is also linear in the number of vertices. To see this, let G be a graph with n vertices and m edges. Suppose that $\text{cr}(G) \leq cm$. If $m < 4n$ then $\text{cr}(G) \leq 4cn$ and we are done. Otherwise $\text{cr}(G) \geq m^3/64n^2$ by the crossing lemma. Thus $m \leq 8\sqrt{cn}$ and $\text{cr}(G) \leq 8c^{3/2}n$.

for every graph G in the family, for some constant c . For example, Pach and Tóth [63] proved that graphs of bounded genus⁵ and bounded degree have linear crossing number. Our main result states that bounded-degree graphs that exclude a fixed graph as a minor⁶ have linear crossing number.

Theorem 1.1. *For every graph H and integer Δ , there is a constant $c = c(H, \Delta)$, such that every H -minor-free graph G with maximum degree at most Δ has crossing number $\text{cr}(G) \leq c|G|$.*

Theorem 1.1 implies the above-mentioned result of Pach and Tóth [63], since graphs of bounded genus exclude a fixed graph as a minor (although the dependence on Δ is different in the two proofs; see Section 6). Moreover, Theorem 1.1 is stronger than the above-mentioned result of Pach and Tóth [63], since there are graphs with a fixed excluded minor and unbounded genus.⁷ For other recent work on minors and crossing number see [16, 17, 18, 38, 40, 44, 45, 55, 64].

Note that the assumption of bounded degree in Theorem 1.1 is unavoidable. For example, $K_{3,n}$ has no K_5 -minor, yet has $\Omega(n^2)$ crossing number [69, 54]. Conversely, bounded degree does not by itself guarantee linear crossing number. For example, a random cubic graph on n vertices has $\Omega(n)$ bisection width [21, 26], which implies that it has $\Omega(n^2)$ crossing number [32, 49].

The proof of Theorem 1.1 is based on *planar decompositions*, which are introduced in Sections 2 and 3. This combinatorial structure generalises tree decompositions by allowing an arbitrary planar graph to index the decomposition. We prove that planar decompositions and the crossing number are intimately related (Section 4). In particular, a graph with bounded degree has linear crossing number if and only if it has a planar decomposition with bounded width and linear order (Theorem 4.5). We study planar decompositions of: K_5 -minor-free graphs (Section 5), graphs embedded in surfaces (Section 6), and finally graphs with an excluded minor (Section 7). One of the main contributions of this paper is to prove that every graph that excludes a fixed graph as a minor has a planar decomposition with bounded width and linear order. Theorem 1.1 easily follows.

1.1. Complementary results. A graph drawing is *rectilinear* (or *geometric*) if each edge is represented by a straight line-segment. The *rectilinear crossing number*

⁵Let \mathbb{S}_γ be the orientable surface with $\gamma \geq 0$ handles. An *embedding* of a graph in \mathbb{S}_γ is a crossing-free drawing in \mathbb{S}_γ . A *2-cell embedding* is an embedding in which each region of the surface (bounded by edges of the graph) is an open disk. The (*orientable*) *genus* of a graph G is the minimum γ such that G has a 2-cell embedding in \mathbb{S}_γ . In what follows, by a *face* we mean the set of vertices on the boundary of the face. Let $F(G)$ be the set of faces in an embedded graph G . See the monograph by Mohar and Thomassen [52] for a thorough treatment of graphs on surfaces.

⁶Let vw be an edge of a graph G . Let G' be the graph obtained by identifying the vertices v and w , deleting loops, and replacing parallel edges by a single edge. Then G' is obtained from G by *contracting* vw . A graph H is a *minor* of a graph G if H can be obtained from a subgraph of G by contracting edges. A family of graphs \mathcal{F} is *minor-closed* if $G \in \mathcal{F}$ implies that every minor of G is in \mathcal{F} . \mathcal{F} is *proper* if it is not the family of all graphs. A deep theorem of Robertson and Seymour [78] states that every proper minor-closed family can be characterised by a finite family of excluded minors. Every proper minor-closed family is a subset of the H -minor-free graphs for some graph H . We thus focus on minor-closed families with one excluded minor.

⁷For example, the complete bipartite graph $K_{3,n}$ has no K_5 -minor and has unbounded genus [72]. Seese and Wessel [81] constructed a family of graphs, each with no K_8 -minor and maximum degree 5, and with unbounded genus.

of a graph G , denoted by $\overline{\text{cr}}(G)$, is the minimum number of crossings in a rectilinear drawing of G ; see [3, 10, 20, 51, 71, 79]. A rectilinear drawing is *convex* if the vertices are positioned on a circle. The *convex* (or *outerplanar*, *circular*, or *1-page book*) *crossing number* of a graph G , denoted by $\text{cr}^*(G)$, is the minimum number of crossings in a convex drawing of G ; see [22, 74, 83]. Obviously

$$\text{cr}(G) \leq \overline{\text{cr}}(G) \leq \text{cr}^*(G)$$

for every graph G . *Linear* rectilinear and *linear* convex crossing numbers are defined in an analogous way to linear crossing number.

It is unknown whether an analogue of Theorem 1.1 holds for rectilinear crossing number.⁸ On the other hand, we prove that $K_{3,3}$ -minor-free graphs with bounded degree have linear rectilinear crossing number (Section 9).

Theorem 1.2. *For every integer Δ , there is a constant $c = c(\Delta)$, such that every $K_{3,3}$ -minor-free graph G with maximum degree at most Δ has rectilinear crossing number $\overline{\text{cr}}(G) \leq c|G|$.*

An analogue of Theorem 1.1 for convex crossing number does not hold, even for planar graphs, since Shahrokhi et al. [83] proved that the $n \times n$ planar grid G_n (which has maximum degree 4) has convex crossing number $\Omega(|G_n| \log |G_n|)$. It is natural to ask which property of the planar grid forces up the convex crossing number. In some sense, we show that tree-width⁹ is one answer to this question. In particular, G_n has tree-width n . More generally, we prove that every graph with large tree-width has many crossings on some edge in every convex drawing (Proposition 8.5). On the other hand, we prove that graphs with bounded tree-width and bounded degree have linear convex crossing number (Section 8).

Theorem 1.3. *For all integers k and Δ , there is a constant $c = c(k, \Delta)$, such that every graph G with tree-width at most k and maximum degree at most Δ has convex crossing number $\overline{\text{cr}}(G) \leq c|G|$.*

Again, the assumption of bounded degree in Theorem 1.3 is unavoidable since $K_{3,n}$ has tree-width 3.

2. Graph decompositions

Let G and D be graphs, such that each vertex of D is a set of vertices of G (called a *bag*). Note that we allow distinct vertices of D to be the same set of vertices in G ; that is, $V(D)$ is a multiset. For each vertex v of G , let $D(v)$ be the subgraph of D induced by the bags that contain v . Then D is a *decomposition* of G if:

- $D(v)$ is connected and nonempty for each vertex v of G .
- $D(v)$ and $D(w)$ touch¹⁰ for each edge vw of G .

⁸The crossing number and rectilinear crossing number are not related in general. In particular, for every integer $k \geq 4$, Bienstock and Dean [10] constructed a graph G_k with crossing number 4 and rectilinear crossing number k . It is easily seen that G_k has no K_{14} -minor. However, the maximum degree of G_k increases with k . Thus G_k is not a counterexample to an analogue of Theorem 1.1 for rectilinear crossing number.

⁹Tree-width is a minor-monotone parameter that is defined in Section 2.

¹⁰Let A and B be subgraphs of a graph G . Then A and B *intersect* if $V(A) \cap V(B) \neq \emptyset$, and A and B *touch* if they intersect or $v \in V(A)$ and $w \in V(B)$ for some edge vw of G .

Decompositions, when D is a tree, were introduced by Robertson and Seymour [76]. Diestel and Kühn [27]¹¹ first generalised the definition for arbitrary graphs D .

Let D be a decomposition of a graph G . The *width* of D is the maximum cardinality of a bag. The number of bags that contain a vertex v of G is the *spread* of v in D . The *spread* of D is the maximum spread of a vertex of G . The *order* of D is the number of bags. D has *linear order* if $|D| \leq c|G|$ for some constant c . If the graph D is a tree, then the decomposition D is a *tree decomposition*. If the graph D is a cycle, then the decomposition D is a *cycle decomposition*. The decomposition D is *planar* if the graph D is planar. The *genus* of the decomposition D is the genus of the graph D .

Note that decompositions generalise minors since D is a decomposition of G with width 1 if and only if G is a minor of D .

A decomposition D of a graph G is *strong* if $D(v)$ and $D(w)$ intersect for each edge vw of G . The *tree-width* of G , denoted by $\text{tw}(G)$, is 1 less than the minimum width of a strong tree decomposition of G . For example, a graph has tree-width 1 if and only if it is a forest. Graphs with tree-width 2 (called *series-parallel*) are planar, and are characterised as those graphs with no K_4 -minor. Tree-width is particularly important in structural and algorithmic graph theory; see the surveys [12, 68].

Tree decompositions are not powerful enough for applications to crossing number: even the $n \times n$ planar grid has tree-width n . We show in Section 4 that planar decompositions are the right type of decomposition for applications to crossing number. It is tempting to define the ‘planar-width’ of a graph G to be the minimum width in a planar decomposition of G . However, by the following lemma of Diestel and Kühn [27], every graph would then have bounded planar-width. We include the proof for completeness.

Lemma 2.1 ([27]). *Every graph G has a strong planar decomposition of width 2, spread $|G|$, and order $\binom{|G|+1}{2}$.*

Proof. Let $n := |G|$ and say $V(G) = \{1, 2, \dots, n\}$. Define a graph D with vertex set $V(D) := \{\{i, j\} : 1 \leq i \leq j \leq n\}$ and edge set

$$E(D) := \{\{i, j\}\{i + 1, j\} : 1 \leq i \leq n - 1, 1 \leq j \leq n\}.$$

Then D is a planar subgraph of the $n \times n$ grid; see Figure 1. For each vertex i of G , the set of bags that contain i is $\{\{i, j\} : 1 \leq j \leq n\}$, which induces a (connected) n -vertex path in D . For each edge ij of G , the bag $\{i, j\}$ contains i and j . Therefore D is a strong decomposition of G . The width is 2, since each bag has at most two vertices. Each vertex is in n bags. \square

¹¹A decomposition was called a *connected decomposition* by Diestel and Kühn [27]. Similar definitions were introduced by Agnew [2]. Decompositions can also be defined in terms of the lexicographic product. For a graph G and integer $k \geq 1$, the lexicographic product $G \cdot K_k$ is the graph with vertex set $V(G) \times [k]$, where $(v, i)(w, j)$ is an edge of $G \cdot K_k$ if and only if $vw \in E(G)$, or $v = w$ and $i \neq j$. That is, each vertex of G is ‘blown up’ by a copy of K_k , and each edge of G is ‘blown up’ by a copy of $K_{k,k}$. It is easily seen that D is a decomposition of G with width k if and only if G is a minor of $D \cdot K_k$. With this viewpoint, a similar result to Lemma 2.1 was obtained by Nešetřil and Ossona De Mendez [56], who observed that every graph is a minor of $D \cdot K_2$ for some planar graph D .




FIGURE 1. A strong planar decomposition of K_6 with width 2 and order 21; the subgraph $D(3)$ is highlighted.

The planar decomposition in Lemma 2.1 has large order (quadratic in $|G|$). The remainder of this paper focuses on planar decompositions with linear order.

Strong tree decompositions are the most widely studied decompositions in the literature. This paper focuses on decompositions that are not necessarily strong. One advantage is that every graph obviously has a decomposition isomorphic to itself (with width 1). On the other hand, if G has a strong decomposition D of width k , then

$$(1) \quad \|G\| \leq \binom{k}{2} |D|.$$

It follows that if G has a strong decomposition isomorphic to itself then the width is at least $\sqrt{\frac{2\|G\|}{|G|}}$, which is unbounded for dense graphs, as observed by Diestel and Kühn [27]. Note that if G has a (nonstrong) decomposition D of width k , then

$$(2) \quad \|G\| \leq k^2 \|D\| + \binom{k}{2} |D|.$$

Every tree T satisfies the Helly property: every collection of pairwise intersecting subtrees of T have a vertex in common. It follows that if a tree T is a strong decomposition of G then every clique¹² of G is contained in some bag of T . Other graphs do not have this property. It will be desirable (for performing k -sums in Section 3) that (nontree) decompositions have a similar property. We therefore introduce the following definitions.

For $p \geq 0$, a p -clique is a clique of cardinality p . A $(\leq p)$ -clique is a clique of cardinality at most p . For $p \geq 2$, a decomposition D of a graph G is a p -decomposition if each $(\leq p)$ -clique of G is a subset of some bag of D , or is a subset of the union of two adjacent bags of D . An $\omega(G)$ -decomposition of G is called an ω -decomposition. A p -decomposition D of G is *strong* if each $(\leq p)$ -clique of G is a subset of some bag of D . Observe that a (strong) 2-decomposition is the same as a (strong) decomposition, and a (strong) p -decomposition also is a (strong) q -decomposition for all $q \in [2, p]$.

¹²A *clique* of a graph G is a set of pairwise adjacent vertices in G . The maximum cardinality of a clique of G is denoted by $\omega(G)$.

3. Manipulating decompositions

In this section we describe four tools for manipulating graph decompositions that are repeatedly used in the remainder of the paper.

Tool #1. Contracting a Decomposition: Our first tool describes the effect of contracting an edge in a decomposition.

Lemma 3.1 (Contraction). *Suppose that D is a planar (strong) p -decomposition of a graph G with width k . Say XY is an edge of D . Then the decomposition D' obtained by contracting the edge XY into the vertex $X \cup Y$ is a planar (strong) p -decomposition of G with width $\max\{k, |X \cup Y|\}$. In particular, if $|X \cup Y| \leq k$ then D' also has width k .*

Proof. Contracting edges preserves planarity. Thus D' is planar. Contracting edges preserves connectiveness. Thus $D'(v)$ is connected for each vertex v of G . Contracting the edge XY obviously maintains the required properties for each $(\leq p)$ -clique of G . \square

Lemma 3.1 can be used to decrease the order of a decomposition at the expense of increasing the width. The following observation is a corollary of Lemma 3.1.

Corollary 3.2. *Suppose that D is a (strong) p -decomposition of a graph G with width k , and that D has a matching¹³ M . The decomposition obtained from D by contracting M is a (strong) p -decomposition of G with width at most $2k$ and order $|D| - |M|$.* \square

Lemma 3.3. *Suppose that a graph G has a (strong) planar p -decomposition D of width k and order at most $c|G|$ for some $c \geq 1$. Then G has a (strong) planar p -decomposition of width $c'k$ and order $|G|$, for some c' depending only on c .*

Proof. Without loss of generality, D is a planar triangulation. Biedl et al. [9] proved that every planar triangulation on n vertices has a matching of at least $\frac{n}{3}$ edges. Applying this result to D , and by Corollary 3.2, G has a (strong) planar p -decomposition of width at most $2k$ and order at most $\frac{2}{3}|D|$. By induction, for every integer $i \geq 1$, G has a (strong) planar p -decomposition of width $2^i k$ and order at most $\max\{(\frac{2}{3})^i |D|, 1\}$. With $i := \lceil \log_{3/2} c \rceil$, the assumption that $|D| = c|G|$ implies that G has a (strong) planar p -decomposition of width $2^i k$ and order $|G|$. \square

Tool #2. Composing Decompositions: Our second tool describes how two decompositions can be composed.

Lemma 3.4 (Composition Lemma). *Suppose that D is a (strong) p -decomposition of a graph G with width k , and that J is a decomposition of D with width ℓ . Then G has a (strong) p -decomposition isomorphic to J with width $k\ell$.*

Proof. Let J' be the graph isomorphic to J that is obtained by renaming each bag $Y \in V(J)$ by $Y' := \{v \in V(G) : v \in X \in Y \text{ for some } X \in V(D)\}$. There are at most ℓ vertices $X \in Y$, and at most k vertices $v \in X$. Thus each bag of J' has at most $k\ell$ vertices.

First we prove that $J'(v)$ is connected for each vertex v of G . Let A' and B' be two bags of J' that contain v . Let A and B be the corresponding bags in D . Thus

¹³A *matching* is a set of pairwise disjoint edges.

$v \in X_1$ and $v \in X_t$ for some bags $X_1, X_t \in V(D)$ such that $X_1 \in A$ and $X_t \in B$ (by the construction of J'). Since $D(v)$ is connected, there is a path X_1, X_2, \dots, X_t in D such that v is in each X_i . In particular, each $X_i X_{i+1}$ is an edge of D . Now $J(X_i)$ and $J(X_{i+1})$ touch in J . Thus there is path in J between any vertex of J that contains X_1 and any vertex of J that contains X_t , such that every bag in the path contains some X_i . In particular, there is a path P in J between A and B such that every bag in P contains some X_i . Let $P' := \{Y' : Y \in P\}$. Then $v \in Y'$ for each bag Y' of P' (by the construction of J'). Thus P' is a connected subgraph of J' that includes A' and B' , and v is in every such bag. Therefore $J'(v)$ is connected.

It remains to prove that for each $(\leq p)$ -clique C of G ,

- (a) C is a subset of some bag of J' , or
- (b) C is a subset of the union of two adjacent bags of J' .

Moreover, we must prove that if D is strong then case (a) always occurs. Since D is a p -decomposition of G ,

- (1) $C \subseteq X$ for some bag $X \in V(D)$, or
- (2) $C \subseteq X_1 \cup X_2$ for some edge $X_1 X_2$ of D .

First suppose that case (1) applies, which always occurs if D is strong. Then there is some bag $Y \in V(J)$ such that $X \in Y$ (since X is a vertex of D and J is a decomposition of D). Thus $C \subseteq Y'$ by the construction of J' . Now suppose that case (2) applies. Then $D(X_1)$ and $D(X_2)$ touch in D . That is, X_1 and X_2 are in a common bag of D , or $X_1 \in Y_1$ and $X_2 \in Y_2$ for some edge $Y_1 Y_2$ of D . If X_1 and X_2 are in a common bag Y , then since $C \subseteq X_1 \cup X_2$, we have $C \subseteq Y'$ by the construction of J' ; that is, case (a) occurs. Otherwise, $X_1 \in Y_1$ and $X_2 \in Y_2$ for some edge $Y_1 Y_2$ of D . Then $C \cap X_1 \subseteq Y_1'$ and $C \cap X_2 \subseteq Y_2'$. Since $C \subseteq X_1 \cup X_2$ we have $C \subseteq Y_1' \cup Y_2'$; that is, case (b) occurs. \square

Tool #3. ω -Decompositions: The third tool converts a decomposition into an ω -decomposition with a small increase in the width. A graph G is d -degenerate if every subgraph of G has a vertex of degree at most d .

Lemma 3.5. *Every d -degenerate graph G has a strong ω -decomposition isomorphic to G of width at most $d + 1$.*

Proof. It is well-known (and easily proved) that G has an acyclic orientation¹⁴ such that each vertex has indegree at most d . Replace each vertex v by the bag $\{v\} \cup N_G^-(v)$. Every subgraph of G has a *sink*. Thus every clique is a subset of some bag. The set of bags that contain a vertex v are indexed by $\{v\} \cup N_G^+(v)$, which induces a connected subgraph in G . Thus we have a strong ω -decomposition. Each bag has cardinality at most $d + 1$. \square

Lemma 3.6. *Suppose that D is a decomposition of a d -degenerate graph G of width k . Then G has a strong ω -decomposition isomorphic to D of width $k(d + 1)$.*

¹⁴If each edge of a graph G is directed from one endpoint to the other, then we speak of an *orientation* of G with arc set $A(G)$. An orientation with no directed cycle is *acyclic*. For each vertex v of an orientation of G , let $N_G^-(v) := \{w \in V(G) : (w, v) \in A(G)\}$ and $N_G^+(v) := \{w \in V(G) : (v, w) \in A(G)\}$. The *indegree* and *outdegree* of v are $|N_G^-(v)|$ and $|N_G^+(v)|$ respectively. A *sink* is a vertex with outdegree 0.

Proof. By Lemma 3.5, G has a strong ω -decomposition isomorphic to G of width d . By Lemma 3.4, G has a strong ω -decomposition isomorphic to D with width $k(d + 1)$. \square

In Lemma 3.6, the ‘blow-up’ in the width is bounded by a constant factor for the graphs that we are interested in: even in the most general setting, H -minor-free graphs are $c|H|\sqrt{\log |H|}$ -degenerate for some constant c [47, 89, 90].

Tool #4. Clique-Sums of Decompositions: Our fourth tool describes how to determine a planar decomposition of a clique-sum of two graphs, given planar decompositions of the summands.¹⁵ Let G_1 and G_2 be disjoint graphs. Suppose that C_1 and C_2 are k -cliques of G_1 and G_2 respectively, for some integer $k \geq 0$. Let $C_1 = \{v_1, v_2, \dots, v_k\}$ and $C_2 = \{w_1, w_2, \dots, w_k\}$. Let G be a graph obtained from $G_1 \cup G_2$ by identifying v_i and w_i for each $i \in [1, k]$, and possibly deleting some of the edges $v_i v_j$. Then G is a k -sum of G_1 and G_2 joined at $C_1 = C_2$. An ℓ -sum for some $\ell \leq k$ is called a $(\leq k)$ -sum. For example, if G_1 and G_2 are planar then it is easily seen that every (≤ 2) -sum of G_1 and G_2 is also planar, as illustrated in Figure 2.




FIGURE 2. Clique-sums of planar graphs. In (b) we can assume that the edges $v_1 v_2$ and $w_1 w_2$ are respectively on the outerfaces of G_1 and G_2 .

Lemma 3.7 (Clique Sums). *Suppose that for integers $p \leq q$, a graph G is a $(\leq p)$ -sum of graphs G_1 and G_2 , and each G_i has a (strong) planar q -decomposition D_i of width k_i . Then G has a (strong) planar q -decomposition of width $\max\{k_1, k_2\}$ and order $|D_1| + |D_2|$.*

Proof. Let $C := V(G_1) \cap V(G_2)$. Then C is a $(\leq p)$ -clique, and thus a $(\leq q)$ -clique, of both G_1 and G_2 . Thus for each i ,

- (1) $C \subseteq X_i$ for some bag X_i of D_i , or
- (2) $C \subseteq X_i \cup Y_i$ for some edge $X_i Y_i$ of D_i .

If (1) is applicable, which is the case if D_i is strong, then consider $Y_i := X_i$ in what follows.

Let D be the graph obtained from the disjoint union of D_1 and D_2 by adding edges $X_1 X_2$, $X_1 Y_2$, $Y_1 X_2$, and $Y_1 Y_2$. By considering $X_1 Y_1$ to be on the outerface of G_1 and $X_2 Y_2$ to be on the outerface of G_2 , observe that D is planar, as illustrated in Figure 3.

¹⁵Leaños and Salazar [48] recently proved some related results on the additivity of crossing numbers.




FIGURE 3. Sum of (a) strong planar decompositions, (b) planar decompositions.

We now prove that $D(v)$ is connected for each vertex v of G . If $v \notin V(G_1)$ then $D(v) = D_2(v)$, which is connected. If $v \notin V(G_2)$ then $D(v) = D_1(v)$, which is connected. Otherwise, $v \in C$. Thus $D(v) = D_1(v) \cup D_2(v)$. Since $v \in X_1 \cup Y_1$ and $v \in X_2 \cup Y_2$, and X_1, Y_1, X_2, Y_2 induce a connected subgraph ($\subseteq K_4$) in D , we have that $D(v)$ is connected.

Each $(\leq q)$ -clique B of G is a $(\leq q)$ -clique of G_1 or G_2 . Thus B is a subset of some bag of D , or B is a subset of the union of two adjacent bags of D . Moreover, if D_1 and D_2 are both strong, then B is a subset of some bag of D . Therefore D is a q -decomposition of G , and if D_1 and D_2 are both strong then D is also strong. The width and order of D are obviously as claimed. \square

A graph can be obtained by repeated (≤ 1) -sums of its biconnected components. Thus Lemma 3.7 with $p = 1$ implies:

Corollary 3.8. *Let G be a graph with biconnected components G_1, G_2, \dots, G_t . Suppose that each G_i has a (strong) planar q -decomposition of width k_i and order n_i . Then G has a (strong) planar q -decomposition of width $\max_i k_i$ and order $\sum_i n_i$. \square*

Note that in the proof of Lemma 3.7, if $X_1 \subseteq X_2$ (for example) then, by Lemma 3.1, we can contract the edge X_1X_2 in D and merge the corresponding bags. The width is unchanged and the order is decreased by 1. This idea is repeatedly used in the remainder of the paper.

4. Planar decompositions and the crossing number

The following lemma is the key link between planar decompositions and the crossing number of a graph.

Lemma 4.1. *Suppose that D is a planar decomposition of a graph G with width k . Then the crossing number of G satisfies*

$$\text{cr}(G) \leq 2 \Delta(G)^2 \sum_{X \in V(D)} \binom{|X|+1}{2} \leq k(k+1) \Delta(G)^2 |D|.$$

Moreover, if $s(v)$ is the spread of each vertex v of G in D , then G has a drawing with the claimed number of crossings, in which each edge vw is represented by a polyline with at most $s(v) + s(w) - 2$ bends.

Proof. By the Fáry–Wagner Theorem [37, 92], D has a rectilinear drawing with no crossings. Let $R_\epsilon(X)$ be the open disc of radius $\epsilon > 0$ centred at each vertex X in the drawing of D . For each edge XY of D , let $R_\epsilon(XY)$ be the union of all segments with one endpoint in $R_\epsilon(X)$ and one endpoint in $R_\epsilon(Y)$. For some $\epsilon > 0$,

- (a) $R_\epsilon(X) \cap R_\epsilon(Y) = \emptyset$ for all distinct bags X and Y of D , and
- (b) $R_\epsilon(XY) \cap R_\epsilon(AB) = \emptyset$ for all edges XY and AB of D that have no endpoint in common.

For each vertex v of G , choose a bag S_v of D that contains v . For each vertex v of G , choose a point $p(v) \in R_\epsilon(S_v)$, and for each bag X of D , choose a set $P(X)$ of $\sum_{v \in X} \deg_G(v)$ points in $R_\epsilon(X)$, so that if

$$\mathcal{P} = \cup\{P(X) : X \in V(D)\} \cup \{p(v) : v \in V(G)\}$$

then:

- (c) no two points in \mathcal{P} coincide,
- (d) no three points in \mathcal{P} are collinear, and
- (e) no three segments, each connecting two points in \mathcal{P} , cross at a common point.

The set \mathcal{P} can be chosen iteratively since each disc $R_\epsilon(X)$ is 2-dimensional,¹⁶ but the set of excluded points is 1-dimensional.

Draw each vertex v at $p(v)$. For each edge vw of G , a simple polyline

$$L(vw) = (p(v), x_1, x_2, \dots, x_a, y_1, y_2, \dots, y_b, p(w))$$

(defined by its endpoints and bends) is a *feasible* representation of vw if:

- (1) $a \in [0, s(v) - 1]$ and $b \in [0, s(w) - 1]$,
- (2) each bend x_i is in $P(X_i)$ for some bag X_i containing v ,
- (3) each bend y_i is in $P(Y_i)$ for some bag Y_i containing w ,
- (4) the bags $S_v, X_1, X_2, \dots, X_a, Y_1, Y_2, \dots, Y_b, S_w$ are distinct (unless $S_v = S_w$ in which case $a = b = 0$), and
- (5) consecutive bends in $L(vw)$ occur in adjacent bags of D .

Since $D(v)$ and $D(w)$ touch, there is a feasible polyline that represents vw .

A drawing of G is *feasible* if every edge of G is represented by a feasible polyline, and no two bends coincide. Since each $|P(X)| = \sum_{v \in X} \deg(v)$, there is a feasible drawing. In particular, no edge passes through a vertex by properties (c)–(e), and no three edges have a common crossing point by property (e). By property (1), each edge vw has at most $s(v) + s(w) - 2$ bends.

Now choose a feasible drawing that minimises the total (Euclidean) length of the edges (with $\{p(v) : v \in V(G)\}$ and $\{P(X) : X \in V(D)\}$ fixed).

By properties (a), (b) and (2)–(5), each segment in a feasible drawing is contained within $R_\epsilon(X)$ for some bag X of D , or within $R_\epsilon(XY)$ for some edge XY of D . Consider a crossing in G between edges vw and xy . Since D is drawn without crossings, the crossing point is contained within $R_\epsilon(X)$ for some bag X of D , or within $R_\epsilon(XY)$ for some edge XY of D . Thus some endpoint of vw , say v , and some endpoint of xy , say x , are in a common bag X . In this case, charge the

¹⁶Let Q be a nonempty set of points in the plane. Then Q is *2-dimensional* if it contains a disk of positive radius; Q is *1-dimensional* if it is not 2-dimensional but contains a finite curve; otherwise Q is *0-dimensional*.

crossing to the 5-tuple (vw, v, xy, x, X) . Observe that the number of such 5-tuples is

$$\sum_{X \in V(D)} \sum_{v, x \in X} \deg_G(v) \cdot \deg_G(x).$$

At most four crossings are charged to each 5-tuple (vw, v, xy, x, X) , since by property (4), each of vw and xy have at most two segments that intersect $R_\epsilon(X)$ (which might pairwise cross). We claim that, in fact, at most two crossings are charged to each such 5-tuple.

Suppose on the contrary that at least three crossings are charged to some 5-tuple (vw, v, xy, x, X) . Then two segments of vw intersect $R_\epsilon(X)$ and two segments of xy intersect $R_\epsilon(X)$. In particular, $p(v) \notin R_\epsilon(X)$ and $p(x) \notin R_\epsilon(X)$, and vw and xy each have a bend in $R_\epsilon(X)$. Let (r_1, r_2, r_3) be the 2-segment polyline in the representation of vw , where r_2 is the bend of vw in $R_\epsilon(X)$. Let (t_1, t_2, t_3) be the 2-segment polyline in the representation of xy , where t_2 is the bend of xy in $R_\epsilon(X)$. Since at least three crossings are charged to (vw, v, xy, x, X) , in the set of segments $\{r_1r_2, r_2r_3, t_1t_2, t_2t_3\}$, at most one pair of segments, one from vw and one from xy , do not cross. Without loss of generality, t_1t_2 and r_2r_3 are this pair. Observe that the crossing segments r_1r_2 and t_1t_2 are the diagonals of the convex quadrilateral $r_1t_2r_2t_1$. Replace the segments r_1r_2 and t_1t_2 by the segments r_1t_2 and t_1r_2 , which are on opposite sides of the quadrilateral. Thus the combined length of r_1t_2 and t_1r_2 is less than the combined length of r_1r_2 and t_1t_2 . Similarly, replace the segments r_3r_2 and t_3t_2 by the segments r_3t_2 and t_3r_2 . We obtain a feasible drawing of G with less total length. This contradiction proves that at most two crossings are charged to each 5-tuple (vw, v, xy, x, X) .

Thus the number of crossings is at most twice the number of 5-tuples. Therefore the number of crossings is at most

$$2 \sum_{X \in V(D)} \sum_{v, x \in X} \deg_G(v) \cdot \deg_G(x) \leq 2 \Delta(G)^2 \sum_{X \in V(D)} \binom{|X|+1}{2}. \quad \square$$




FIGURE 4. In the proof of Lemma 4.1, to shorten the total edge length the crossed segments are replaced by the dashed segments.

Note that the bound on the crossing number in Lemma 4.1 is within a constant factor of optimal for the complete graph. An easy generalisation of Lemma 2.1

proves that for all $n \geq k \geq 2$, K_n has a strong planar decomposition of width k and order at most $c(\frac{n}{k})^2$ for some constant c . Thus Lemma 4.1 implies that $\text{cr}(K_n) \leq ck(k+1) \Delta(K_n)^2 (\frac{n}{k})^2 \leq cn^4$, which is within a constant factor of optimal [71].

The following result is converse to Lemma 4.1.

Lemma 4.2. *Let G be a graph with n isolated vertices. Suppose that G has a drawing with c crossings in which q nonisolated vertices are not incident to a crossed edge. Then G has a planar decomposition of width 2 and order $\lceil \frac{n}{2} \rceil + q + c$, and G has a strong planar decomposition of width 2 and order $\lceil \frac{n}{2} \rceil + c + \|G\|$.*

Proof. First, pair the isolated vertices of G . Each pair can form one bag in a decomposition of width 2, adding $\lceil \frac{n}{2} \rceil$ to the order. Now assume that G has no isolated vertices.

We first construct the (nonstrong) decomposition. Arbitrarily orient each edge of G . Let D be the planar graph obtained from the given drawing of G by replacing each vertex v by the bag $\{v\}$, and replacing each crossing between arcs (v, w) and (x, y) of G by a degree-4 vertex $\{v, x\}$. Thus an arc (v, w) of G is replaced by some path $\{v\}\{v, x_1\}\{v, x_2\} \dots \{v, x_r\}\{w\}$ in D . In particular, $D(v)$ and $D(w)$ touch at the edge $\{v, x_r\}\{w\}$. Moreover, $D(v)$ is a (connected) tree for each vertex v of G . Thus D is a decomposition of G . Each bag contains at most two vertices. The order is $|G| + c$. For each vertex v of G that is incident to some crossed edge, $\{v\}\{v, x\}$ is an edge in D for some vertex x . Contract the edge $\{v\}\{v, x\}$ in D and merge the bags. By Lemma 3.1, D remains a planar decomposition of width 2. The order is now $q + c$.

Now we make D strong. For each edge vw of G , there is an edge XY of D where $D(v)$ and $D(w)$ touch. That is, $v \in X$ and $w \in Y$. Replace XY by the path $X\{v, w\}Y$. Now each edge of G is in some bag of D , and D is strong. This operation introduces a further $\|G\|$ bags. Thus the order is $q + c + \|G\|$. For each nonisolated vertex v that is not incident to a crossed edge, choose an edge vw incident to v . Then $\{v\}\{v, w\}$ is an edge in D ; contract this edge and merge the bags. By Lemma 3.1, D remains a strong decomposition of G with width 2. The order is now $c + \|G\|$. □

The following two special cases of Lemma 4.2 are of particular interest.

Corollary 4.3. *Every graph G has a planar decomposition of width 2 and order $|G| + \text{cr}(G)$. Every graph G has a strong planar decomposition of width 2 and order $|G| + \|G\| + \text{cr}(G)$.* □

Corollary 4.4. *Every planar graph G with n isolated vertices has a strong planar decomposition of width 2 and order $\lceil \frac{n}{2} \rceil + \|G\| \leq 3|G| - 6$.* □

The relationship between planar decompositions and graphs with linear crossing number is summarised as follows.

Theorem 4.5. *The following are equivalent for a graph G of bounded degree:*

- (1) $\text{cr}(G) \leq c_1|G|$ for some constant c_1 .
- (2) G has a planar decomposition with width c_2 and order $|G|$ for some constant c_2 .
- (3) G has a planar decomposition with width 2 and order $c_3|G|$ for some constant c_3 .

Proof. Lemma 4.2 implies that (1) \Rightarrow (3). Lemma 3.3 implies that (3) \Rightarrow (2). Lemma 4.1 implies that (2) \Rightarrow (1). \square

5. K_5 -minor-free graphs

In this section we prove the following upper bound on the crossing number.

Theorem 5.1. *Every K_5 -minor-free graph G has crossing number*

$$\text{cr}(G) < \frac{20}{3} \Delta(G)^2 |G|.$$

The proof of Theorem 5.1 is based on Theorem 5.2 below, in which we construct ω -decompositions of K_5 -minor-free graphs G . Since $\omega(G) \leq 4$ and each clique can be spread over two bags, it is natural to consider ω -decompositions of width 2.

Theorem 5.2. *Every K_5 -minor-free graph G with $|G| \geq 3$ has a planar ω -decomposition of width 2 and order at most $\frac{4}{3}|G| - 2$, with at most $|G| - 2$ bags of cardinality 2, and at most $\frac{1}{3}|G|$ bags of cardinality 1.*

The proof of Theorem 5.2 is based on the following classical theorem of Wagner [93] and the two following lemmas. Here V_8 is the the 8-vertex Möbius ladder, which is the graph obtained from the 8-cycle by adding an edge between each pair of antipodal vertices; see Figure 5(a).

Theorem 5.3 ([93]). *A graph is K_5 -minor-free if and only if it can be obtained from planar graphs and V_8 by (≤ 3) -sums.*

Lemma 5.4. *Every K_5 -minor-free graph G with $|G| \geq 3$ has a partition of $E(G)$ into three sets E^1, E^2, E^3 such that:*

- *Each of E^1, E^2, E^3 has at most $|G| - 2$ edges.*
- *Every triangle has one edge in each of E^1, E^2, E^3 .*
- *If a subgraph H of G is isomorphic to V_8 , then $E(H) \cap E^j$ is a perfect matching in H for all j .*

Moreover, if G is edge-maximal (with no K_5 -minor), then every vertex is incident to an edge in E^j and an edge in E^ℓ for some $j \neq \ell$.

Proof. By Theorem 5.3, we need only consider the following three cases.

Case (a). G is planar: Let G' be a planar triangulation of G . By the Four-Colour Theorem [75], G' has a proper vertex-colouring with colours a, b, c, d . Now determine a Tait edge-colouring [88]. Let E^1 be the set of edges of G' whose endpoints are coloured ab or cd . Let E^2 be the set of edges of G' whose endpoints are coloured ac or bd . Let E^3 be the set of edges of G' whose endpoints are coloured ad or bc . Since the vertices of each triangle are 3-coloured, the edges of each triangle are in distinct E^j . In particular, the edges of each face of G' are in distinct E^j . Each edge of G' is in two of the $2|G| - 4$ faces of G' . Thus $|E^j| = |G| - 2$. The sets $E^j \cap E(G)$ thus satisfy the first two properties for G . Since V_8 is nonplanar, G has no V_8 subgraph, and the third property is satisfied vacuously. Finally, if G is edge-maximal, then $G' = G$, each vertex v is in some face, and v is incident to two edges in distinct sets.




FIGURE 5. (a) The graph V_8 . (b) Drawing of V_8 with one crossing. (c) Planar ω -decomposition of V_8 with width 2 and order 7. (d) Strong planar decomposition of V_8 with width 2 and order 12. (e) Strong planar ω -decomposition of V_8 with width 4 and order 4.

Case (b). $G = V_8$: Using the vertex-numbering in Figure 5(a), let

$$\begin{aligned} E^1 &:= \{12, 34, 56, 78\}, \\ E^2 &:= \{23, 45, 67, 81\}, \\ E^3 &:= \{15, 26, 37, 48\}. \end{aligned}$$

Each E^j is a matching of four edges. The claimed properties follow.

Case (c). G is a (≤ 3) -sum of two smaller K_5 -minor-free graphs G_1 and G_2 : Let C be the join set. By induction, there is a partition of each $E(G_i)$ into three sets E_i^1, E_i^2, E_i^3 with the desired properties. Permute the set indices so that for each edge e with endpoints in C , $e \in E_1^j \cap E_2^j$ for some j . This is possible because C is a (≤ 3) -clique in G_1 and G_2 .

For each $j = 1, 2, 3$, let $E^j := E_1^j \cup E_2^j$. If $|C| \leq 2$, then $|E^j| \leq |E_1^j| + |E_2^j| \leq (|G_1| - 2) + (|G_2| - 2) = |G_1| + |G_2| - 4 \leq |G| - 2$, as desired. Otherwise, C is a triangle in G_1 and G_2 , and $|E_1^j \cap E_2^j| = 1$. Thus $|E^j| \leq |E_1^j| + |E_2^j| - 1 \leq (|G_1| - 2) + (|G_2| - 2) - 1 = |G_1| + |G_2| - 5 = |G| - 2$, as desired. Each triangle of G is in G_1 or G_2 , and thus has one edge in each set E^j .

Consider a V_8 subgraph H of G . Since V_8 is edge-maximal K_5 -minor-free, H is an induced subgraph. Since V_8 is 3-connected and triangle-free, H is a subgraph of G_1 or G_2 . Thus $H \cap E^j$ is a perfect matching of H by induction.

If G is edge-maximal, then G_1 and G_2 are both edge-maximal. Thus every vertex v of G is incident to at least two edges in distinct sets (since the same property holds for v in G_1 or G_2). \square

For a set E of edges in a graph G , a vertex v of G is *E-isolated* if v is incident to no edge in E .

Lemma 5.5. *Suppose that E is a set of edges in a K_5 -minor-free graph G such that every triangle of G has exactly one edge in E , and if S is a subgraph of G isomorphic to V_8 then $E(S) \cap E$ is a perfect matching in S . Let V be the set of E -isolated vertices in G . Then G has a planar ω -decomposition D of width 2 with $V(D) = \{\{v\} : v \in V\} \cup \{\{v, w\} : vw \in E\}$ with no duplicate bags.*

Proof. By Theorem 5.3, we need only consider the following four cases.

Case (a). $G = K_4$: Say $V(G) = \{v, w, x, y\}$. Without loss of generality, $E = \{vw, xy\}$. Thus $V = \emptyset$. Then $D := K_2$, with bags $\{v, w\}$ and $\{x, y\}$, is the desired decomposition of G . Now assume that $G \neq K_4$.

Case (b). $G = V_8$: Thus E is a perfect matching of G . Then $D := K_4$, with one bag for each edge in E , is the desired decomposition of G .

Case (c). G is planar and has no separating triangle (see Figure 6): Fix a plane drawing of G . Thus every triangle of G is a face. Initially, let D be the planar decomposition of G with $V(D) := \{\{v\} : v \in V(G)\}$ and $E(D) := \{\{v\}\{w\} : vw \in E(G)\}$. For each edge $vw \in E$, introduce a new bag $\{v, w\}$ in D , and replace the edge $\{v\}\{w\}$ by the path $\{v\}\{v, w\}\{w\}$. Thus D is a planar subdivision of G . Now consider each triangle uvw of G . Without loss of generality, $vw \in E$. Replace the path $\{v\}\{u\}\{w\}$ in D by the edge $\{u\}\{v, w\}$. Since uvw is a face with only one edge in E , D remains planar. Moreover, $D(v)$ is a connected star for each vertex v of G . Since G has no separating triangle and $G \neq K_4$, each clique is a (≤ 3) -clique. Thus, by construction, each clique is contained in a bag of D , or is contained in the union of two adjacent bags of G . Therefore D is a planar ω -decomposition of G with width 2. The order is $|G| + |E|$. For each vertex $v \notin V$, there is an edge incident to v that is in E . Choose such an edge $vw \in E$. Thus $\{v\}\{v, w\}$ is an edge of D . Contract this edge and merge the bags. By Lemma 3.1, D remains a planar ω -decomposition. Now $V(D) = \{\{v\} : v \in V\} \cup \{\{v, w\} : vw \in E\}$.

Case (d). G is a (≤ 3) -sum of two smaller K_5 -minor-free graphs G_1 and G_2 : Let C be the join set. Let $E_1 := E \cap E(G_1)$ and $E_2 := E \cap E(G_2)$. Then every triangle of G_i has exactly one edge in E_i . Let V_i be the set of vertices of G_i that are E_i -isolated. By induction, each G_i has a planar ω -decomposition D_i of width 2 with $V(D_i) := E_i \cup V_i$. By Lemma 3.7 with $p = q = 4$ and $k_1 = k_2 = 2$, G has a planar ω -decomposition D of width 2 with $V(D) := V_1 \cup E_1 \cup V_2 \cup E_2$. Moreover, each bag of D_1 that intersects C is adjacent to each bag of D_2 that intersects C .

If there is a duplicate bag X in D , then one copy of X is from D_1 and the other copy is from D_2 , and X intersects C . Thus the two copies are adjacent. Contract the edge XX in D into the bag X . By Lemma 3.1, D remains a planar ω -decomposition of G with width 2. Now D has no duplicate bags.

Every bag in D is either a vertex or an edge of G . If $\{v, w\}$ is a bag of D , then vw is in $E_1 \cup E_2 = E$. Conversely, if $vw \in E$ then $vw \in E_1$ or $vw \in E_2$. Thus the bags of cardinality 2 in D are in one-to-one correspondence with edges in E .




FIGURE 6. An example of the construction of a planar ω -decomposition in case (c) of Lemma 5.5, where $E = \{13, 15, 56, 36\}$ and $V = \{2, 4\}$.

Suppose there is bag $\{v\}$ in D but $v \notin V$. Then $v \in C$, $v \in V_2$, and $v \notin V_1$ (or symmetrically, $v \in V_1$ and $v \notin V_2$). That is, v is incident to an edge $vw \in E_1$ but v is E_2 -isolated. Now $\{v, w\}$ is a bag in D_1 , and $\{v\}$ is a bag in D_2 . These bags are adjacent in D . Contract the edge $\{v, w\}\{v\}$ in D into the bag $\{v, w\}$. By Lemma 3.1, D remains an ω -decomposition of G with width 2. An analogous argument applies if $v \in V_1$ and $v \notin V_2$. We have proved that if $\{v\}$ is a bag of D , then $v \in V$. Conversely, if $v \in V$ then $v \in V_1$ or $v \in V_2$ (possibly both), and there is a bag $\{v\}$ in D . Thus the singleton bags of D are in one-to-one correspondence with vertices in V .

Therefore $V(D) = \{\{v\} : v \in V\} \cup \{\{v, w\} : vw \in E\}$ with no duplicate bags. \square

Proof of Theorem 5.2. Let H be an edge-maximal K_5 -minor-free graph containing G as a spanning subgraph. By Lemma 5.4, there is a partition of $E(H)$ into three sets E^1, E^2, E^3 with the stated properties. We now construct the desired planar ω -decomposition of H , which is also the desired decomposition of G . Let V^j be the set of E^j -isolated vertices in H . Consider an E^1 -isolated vertex v . Each edge incident to v is in $E^2 \cup E^3$. Since v is incident to (at least) two edges in

distinct sets, v is not E^2 -isolated and is not E^3 -isolated. In general, $V^i \cap V^j = \emptyset$ for distinct i and j .¹⁷ Thus $|V^i| \leq \frac{1}{3}|H|$ for some i . By Lemma 5.5, H has a planar ω -decomposition of width 2, with $|E^i| = |H| - 2$ bags of cardinality 2, and $|V^i| \leq \frac{1}{3}|H|$ bags of cardinality 1. \square

Proof of Theorem 5.1. By Theorem 5.2, G has a planar ω -decomposition D of width 2, with at most $|G| - 2$ bags of cardinality 2, and at most $\frac{1}{3}|G|$ bags of cardinality 1. By Lemma 4.1,

$$\text{cr}(G) \leq 2 \Delta(G)^2 \sum_{X \in \mathcal{V}(D)} \binom{|X|+1}{2} \leq 2 \Delta(G)^2 (3(|G| - 2) + \frac{1}{3}|G|) < \frac{20}{3} \Delta(G)^2 |G|. \quad \square$$

The following two propositions, while not used to prove bounds on the crossing number, are of independent interest. The proofs, which are similar to the proof of Theorem 5.2, are in [96]. First we consider strong planar 3-decompositions of K_5 -minor-free graphs.

Proposition 5.6 ([96]). *Every K_5 -minor-free graph G with $|G| \geq 3$ has a strong planar 3-decomposition of width 3 and order $3|G| - 8$. Moreover, for all $n \geq 3$, there is a planar graph G , such that $|G| = n$ and every strong planar 3-decomposition of G with width 3 has order at least $3|G| - 8$.*

It follows from Euler's Formula and Theorem 5.3 that every K_5 -minor-free graph G has at most $3|G| - 6$ edges, and is thus 5-degenerate (also see Lemma 6.4 below). Thus by Lemma 3.5, G has a strong ω -decomposition isomorphic to G of width 5. Since $\omega(G) \leq 4$, it is natural to consider strong ω -decompositions of width 4.

Proposition 5.7 ([96]). *Every K_5 -minor-free graph G with $|G| \geq 4$ has a strong planar ω -decomposition of width 4 and order at most $\frac{4}{3}|G| - 4$. Moreover, for all $n \geq 1$, there is a planar graph G_n , such that $|G_n| = 3n$ and every strong ω -decomposition of G_n with width 4 has order at least $\frac{7}{6}|G_n| - 3$.*

6. Graphs embedded on a surface

Recall that \mathbb{S}_γ is the orientable surface with γ handles. As illustrated in Figure 7, a *cycle* in \mathbb{S}_γ is a closed curve in the surface. A cycle is *contractible* if it is contractible to a point in the surface. A noncontractible cycle is *separating* if it separates \mathbb{S}_γ into two connected components.

Let G be a graph embedded in \mathbb{S}_γ . A *noose* of G is a cycle C in \mathbb{S}_γ that does not intersect the interior of an edge of G . Let $V(C)$ be the set of vertices of G intersected by C . The *length* of C is $|V(C)|$. Pach and Tóth [63] proved that, for some constant c_γ , the crossing number of every graph G of genus γ satisfies¹⁸

$$(3) \quad \text{cr}(G) \leq c_\gamma \sum_{v \in V(G)} \deg(v)^2 \leq 2c_\gamma \Delta(G) \|G\|.$$

The following lemma is probably well-known.

¹⁷Note that it is possible for V_1, V_2, V_3 to partition $V(H)$; for example, when H is a planar Eulerian triangulation.

¹⁸Recently Djidjev and Vrto [33] greatly improved the constant c_γ in (3), and Böröczky et al. [19] extended the bound to graphs embedded on nonorientable surfaces.




FIGURE 7. Cycles on the double torus.

Lemma 6.1. *Every graph with $n \geq 3$ vertices and genus γ has at most*

$$(\sqrt{3\gamma} + 3)n - 6$$

edges.

Proof. Let G be an edge-maximal graph with n vertices, m edges, and genus γ . Suppose that an embedding of G in \mathbb{S}_γ has f faces. Euler’s Formula states that $m - n - f = 2\gamma - 2$. Since G is edge-maximal, every face is bounded by three edges and each edge is in the boundary of two faces. Thus $m = 3f/2$ and $f = 2m/3$. Hence $m = 3n + 6\gamma - 6$. If $\gamma = 0$ then we are done. Now assume that $\gamma \geq 1$. We need to prove that $3n + 6\gamma \leq (\sqrt{3\gamma} + 3)n$. That is, $6\gamma \leq \sqrt{3\gamma}n$, or equivalently $\gamma \leq n^2/12$, which is true since K_n has genus $\lceil (n - 3)(n - 4)/12 \rceil < n^2/12$; see [73, 52]. \square

Equation (3) and Lemma 6.1 imply that

$$(4) \quad \text{cr}(G) \leq c_\gamma \Delta(G) |G|.$$

By Lemma 4.2, G has a planar decomposition of width 2 and order $c_\gamma \Delta(G) |G|$. We now provide an analogous result without the dependence on $\Delta(G)$, but at the expense of an increased bound on the width.

Theorem 6.2. *Every graph G with genus γ has a planar decomposition of width 2^γ and order $3^\gamma |G|$.*

The key to the proof of Theorem 6.2 is the following lemma, whose proof is inspired by similar ideas of Pach and Tóth [63].

Lemma 6.3. *Let G be a graph with a 2-cell embedding in \mathbb{S}_γ for some $\gamma \geq 1$. Then G has a decomposition of width 2, genus at most $\gamma - 1$, and order $3|G|$.*

Proof. Since $\gamma \geq 1$, \mathbb{S}_γ has a noncontractible nonseparating cycle, which can be continuously deformed into a noncontractible nonseparating noose in G . Let C be a noncontractible nonseparating noose in G of minimum length $k := |V(C)|$. Orient C and let $V(C) := (v_1, v_2, \dots, v_k)$ in the order around C . For each vertex $v_i \in V(C)$, let $E^\ell(v_i)$ and $E^r(v_i)$ respectively be the set of edges incident to v_i that are on the left-hand side and right-hand side of C (with respect to the orientation).

Cut the surface along C , and attach a disk to each side of the cut. Replace each vertex $v_i \in V(C)$ by two vertices v_i^ℓ and v_i^r respectively incident to the edges in $E^\ell(v_i)$ and $E^r(v_i)$. Embed v_i^ℓ on the left-hand side of the cut, and embed v_i^r on the right-hand side of the cut. We obtain a graph G' embedded in a surface of genus at most $\gamma - 1$ (since C is nonseparating).

Let $L := \{v_i^\ell : v \in V(C)\}$ and $R := \{v_i^r : v \in V(C)\}$. By Menger's Theorem, the maximum number of disjoint paths between L and R in G' equals the minimum number of vertices that separate L from R in G' . Let Q be a minimum set of vertices that separate L from R in G' . Then there is a noncontractible nonseparating noose in G that only intersects vertices in Q . (It is nonseparating in G since L and R are identified in G .) Thus $|Q| \geq k$ by the minimality of $|V(C)|$. Hence there exist k disjoint paths P_1, P_2, \dots, P_k between L and R in G' , where the endpoints of P_i are v_i^ℓ and $v_{\sigma(i)}^r$, for some permutation σ of $[1, k]$. In the disc with R on its boundary, draw an edge from each vertex $v_{\sigma(i)}^r$ to v_i^r such that no three edges cross at a single point and every pair of edge cross at most once. Add a new vertex $x_{i,j}$ on each crossing point between edges $v_{\sigma(i)}^r v_i^r$ and $v_{\sigma(j)}^r v_j^r$. Let G'' be the graph obtained. Then G'' is embedded in $S_{\gamma-1}$.

We now make G'' a decomposition of G . Replace v_i^ℓ by $\{v_i\}$ and replace v_i^r by $\{v_i\}$. Replace every other vertex v of G by $\{v\}$. Replace each 'crossing' vertex $x_{i,j}$ by $\{v_i, v_j\}$. Now for each vertex $v_i \in V(C)$, add v_i to each bag on the path P_i from v_i^ℓ to $v_{\sigma(i)}^r$. Thus $G''(v_i)$ is a (connected) path. Clearly $G''(v)$ and $G''(w)$ touch for each edge vw of G . Hence G'' is a decomposition of G with genus at most $\gamma - 1$. Since the paths P_1, P_2, \dots, P_k are pairwise disjoint, the width of the decomposition is 2.

It remains to bound the order of G'' . Let $n := |G|$. Observe that G'' has at most $n + k + \binom{k}{2}$ vertices. One of the paths P_i has at most $\frac{n+k}{k}$ vertices. For ease of counting, add a cycle to G' around R . Consider the path in G' that starts at v_i^ℓ , passes through each vertex in P_i , and then takes the shortest route from $v_{\sigma(i)}^r$ around R back to v_i^r . The distance between $v_{\sigma(i)}^r$ and v_i^r around R is at most $\frac{k}{2}$. This path in G' forms a noncontractible nonseparating noose in G (since if two cycles in a surface cross in exactly one point, then both are noncontractible).

The length of this noose in G is at most $\frac{n+k}{k} - 1 + \frac{k}{2}$ (since v_i^ℓ and v_i^r both appeared in the path). Hence $\frac{n+k}{k} - 1 + \frac{k}{2} \geq k$ by the minimality of $|V(C)|$. Thus $k \leq \sqrt{2n}$. Therefore G'' has at most $n + \sqrt{2n} + \binom{\sqrt{2n}}{2} \leq 3n$ vertices. \square

Proof of Theorem 6.2. We proceed by induction on γ . If $\gamma = 0$ then G is planar, and G itself is a planar decomposition of width $1 = 2^0$ and order $n = 3^0 n$. Otherwise, by Lemma 6.3, G has a decomposition D of width 2, genus $\gamma - 1$, and order $3n$. By induction, D has a planar decomposition of width $2^{\gamma-1}$ and order $3^{\gamma-1}(3n) = 3^\gamma n$. By Lemma 3.4 with $p = k = 2$, and $\ell = 2^{\gamma-1}$, G has a planar decomposition of width $2 \cdot 2^{\gamma-1} = 2^\gamma$ and order $3^\gamma n$. \square

Theorem 6.2 and Lemma 4.1 imply that every graph G with genus γ has crossing number $\text{cr}(G) \leq 12^\gamma \Delta(G)^2 |G|$, which for fixed γ , is weaker than the bound of Pach and Tóth [63] in (4). The advantage of our approach is that it generalises for graphs with an arbitrary excluded minor (and the dependence on γ is much smaller).

We now prove that a graph G embedded on a surface has an ω -decomposition with small width and linear order. To do so, we apply Lemma 3.6, which requires a bound on the degeneracy of G .

Lemma 6.4. *Every graph G of genus γ is $(2\sqrt{3\gamma} + 6)$ -degenerate. If $\sqrt{3\gamma}$ is an integer then G is $(2\sqrt{3\gamma} + 5)$ -degenerate.*

Proof. By Lemma 6.1, G has average degree $\frac{2\|G\|}{|G|} < 2(\sqrt{3\gamma} + 3)$. Thus G has a vertex of degree less than $2\sqrt{3\gamma} + 6$. Moreover, if $\sqrt{3\gamma}$ is an integer, then G has a vertex of degree at most $2\sqrt{3\gamma} + 5$. The result follows since every subgraph of G has genus at most γ . \square

Theorem 6.5. *Every graph G of genus γ has a planar ω -decomposition of width $2^\gamma(2\sqrt{3\gamma} + 7)$ and order $3^\gamma |G|$.*

Proof. By Theorem 6.2, G has a planar decomposition D of width at most 2^γ and order $3^\gamma n$. By Lemma 6.4, G is $(2\sqrt{3\gamma} + 6)$ -degenerate. Thus by Lemma 3.6, G has a planar ω -decomposition isomorphic to D with width $2^\gamma(2\sqrt{3\gamma} + 7)$. \square

7. H -minor-free graphs

For integers $h \geq 1$ and $\gamma \geq 0$, Robertson and Seymour [77] defined a graph G to be h -almost embeddable in \mathbb{S}_γ if G has a set X of at most h vertices such that $G \setminus X$ can be written as $G_0 \cup G_1 \cup \dots \cup G_h$ such that:

- G_0 has an embedding in \mathbb{S}_γ .
- The graphs G_1, G_2, \dots, G_h (called *vortices*) are pairwise disjoint.
- There are faces¹⁹ F_1, F_2, \dots, F_h of the embedding of G_0 in \mathbb{S}_γ , such that each $F_i = V(G_0) \cap V(G_i)$.
- If $F_i = (u_{i,1}, u_{i,2}, \dots, u_{i,|F_i|})$ in clockwise order about the face, then G_i has a strong $|F_i|$ -cycle decomposition Q_i of width h , such that each vertex $u_{i,j}$ is in the j -th bag of Q_i .

The following ‘characterisation’ of H -minor-free graphs is a deep theorem by Robertson and Seymour [77]; see the recent survey by Kawarabayashi and Mohar [46].

Theorem 7.1 (Graph Minor Structure Theorem [77]). *For every graph H there is a positive integer $h = h(H)$, such that every H -minor-free graph G can be obtained by $(\leq h)$ -sums of graphs that are h -almost embeddable in some surface in which H cannot be embedded.*

The following theorem is one of the main contributions of this paper.

Theorem 7.2. *For every graph H there is an integer $k = k(H)$, such that every H -minor-free graph G has a planar ω -decomposition of width k and order $|G|$.*

We prove Theorem 7.2 by a series of lemmas.

Lemma 7.3. *Every graph G that is h -almost embeddable in \mathbb{S}_γ has a planar decomposition of width $h(2^\gamma + 1)$ and order $3^\gamma |G|$.*

¹⁹Recall that we equate a face with the set of vertices on its boundary.

Proof. By Theorem 6.2, G_0 has a planar decomposition D of width at most 2^γ and order $3^\gamma |G_0| \leq 3^\gamma |G|$. We can assume that D is connected. For each vortex G_i , add each vertex in the j -th bag of Q_i to each bag of D that contains $u_{i,j}$. The bags of D now contain at most $2^\gamma h$ vertices. Now add X to every bag. The bags of D now contain at most $(2^\gamma + 1)h$ vertices. For each vertex v that is not in a vortex, $D(v)$ is unchanged by the addition of the vortices, and is thus connected. For each vertex v in a vortex G_i , $D(v)$ is the subgraph of D induced by the bags (in the decomposition of G_0) that contain $u_{i,j}$, where v is in the j -th bag of Q_i . Now $Q_i(v)$ is a connected subgraph of the cycle Q_i , and for each vertex $u_{i,j}$, the subgraphs $G_0(u_{i,j})$ and $G_0(u_{i,j+1})$ touch. Thus $D(v)$ is connected. (This argument is similar to that used in Lemma 3.4.) $D(v)$ is connected for each vertex $v \in X$ since D itself is connected. \square

Lemma 7.4. *For all integers $h \geq 1$ and $\gamma \geq 0$ there is a constant $d = d(h, \gamma)$, such that every graph G that is h -almost embeddable in \mathbb{S}_γ is d -degenerate.*

Proof. If G is h -almost embeddable in \mathbb{S}_γ then every subgraph of G is h -almost embeddable in \mathbb{S}_γ . Thus it suffices to prove that if G has n vertices and m edges, then its average degree $\frac{2m}{n} \leq d$. Say each G_i has m_i edges. G has at most hn edges incident to X . Thus $m \leq hn + \sum_{i=0}^h m_i$. By Lemma 6.1, $m_0 < (\sqrt{3^\gamma} + 3)n$. Now $m_i \leq \binom{h}{2} |F_i|$ by Equation (1) with D an $|F_i|$ -cycle. Since G_1, G_2, \dots, G_h are pairwise disjoint, $\sum_{i=1}^h m_i \leq \binom{h}{2} n$. Thus $m < (h + \sqrt{3^\gamma} + 3 + \binom{h}{2})n$. Taking $d = h(h + 1) + 2\sqrt{3^\gamma} + 6$ we are done. \square

Lemmas 3.6, 7.3 and 7.4 imply:

Corollary 7.5. *For all integers $h \geq 1$ and $\gamma \geq 0$ there is a constant $k = k(h, \gamma) \geq h$, such that every graph G that is h -almost embeddable in \mathbb{S}_γ has a planar ω -decomposition of width k and order $3^\gamma |G|$.* \square

Now we bring in ($\leq h$)-sums.

Lemma 7.6. *For all integers $h \geq 1$ and $\gamma \geq 0$, every graph G that can be obtained by ($\leq h$)-sums of graphs that are h -almost embeddable in \mathbb{S}_γ has a planar ω -decomposition of width k and order $\max\{1, 3^\gamma(h + 1)(|G| - h)\}$, where $k = k(h, \gamma)$ from Corollary 7.5.*

Proof. We proceed by induction on $|G|$. If $|G| \leq h$ then the decomposition of G with all its vertices in a single bag satisfies the claim (since $k \geq h$).

Now assume that $|G| \geq h + 1$. If G is h -almost embeddable in \mathbb{S}_γ , then by Corollary 7.5, G has a planar ω -decomposition of width k and order $3^\gamma |G|$, which, since $|G| \geq h + 1$, is at most $3^\gamma(h + 1)(|G| - h)$, as desired.

Otherwise, G is a ($\leq h$)-sum of graphs G_1 and G_2 , each of which, by induction, has a planar ω -decomposition of width k and order $\max\{1, 3^\gamma(h + 1)(|G_i| - h)\}$. By Lemma 3.7, G has a planar ω -decomposition D of width k and order

$$|D| = \max\{1, 3^\gamma(h + 1)(|G_1| - h)\} + \max\{1, 3^\gamma(h + 1)(|G_2| - h)\}.$$

Without loss of generality, $|G_1| \leq |G_2|$. If $|G_2| \leq h$ then $|D| = 2 \leq 3^\gamma(h + 1)(|G| - h)$, as desired. If $|G_1| \leq h$ and $|G_2| \geq h + 1$, then $|D| = 1 + 3^\gamma(h + 1)(|G_2| - h)$, which, since $|G| \geq |G_2| + 1$, is at most $3^\gamma(h + 1)(|G| - h)$, as desired. Otherwise, both $|G_1| \geq h + 1$ and $|G_2| \geq h + 1$. Thus the order of D is $3^\gamma(h + 1)(|G_1| + |G_2| - 2h) \leq 3^\gamma(h + 1)(|G| - h)$, as desired. \square

Proof of Theorem 7.2. Let $h = h(H)$ from Theorem 7.1. Let \mathbb{S}_γ be the surface in Theorem 7.1 in which H cannot be embedded. If G has no H -minor then, by Theorem 7.1, G can be obtained by $(\leq h)$ -sums of graphs that are h -almost embeddable in \mathbb{S}_γ . By Lemma 7.6, G has a planar ω -decomposition of width k and order $3^\gamma(h + 1)|G|$, where $k = k(h, \gamma)$ from Corollary 7.5. By Lemma 3.3, G has a planar ω -decomposition of width k' and order $|G|$, for some k' only depending on k, γ and h (all of which only depend on H). \square

Theorem 7.2 and Lemma 4.1 imply the following quantitative version of Theorem 1.1.

Corollary 7.7. *For every graph H there is a constant $c = c(H)$, such that every H -minor-free graph G has crossing number at most $c\Delta(G)^2|G|$.* \square

It is an open problem whether the dependence on $\Delta(G)$ in Corollary 7.7 can be reduced from quadratic to linear (even with $H = K_5$). We conjecture the stronger result that every H -minor-free graph G has crossing number

$$\text{cr}(G) \leq c_H \sum_{v \in V(G)} \deg(v)^2.$$

Pach and Tóth [63] proved this conjecture for graphs of bounded genus.

8. Graph partitions

A *partition* of a graph is a proper partition of its vertex set. Each part of the partition is called a *bag*. The *width* of partition is the maximum number of vertices in a bag. The *pattern* (or *quotient graph*) of a partition is the graph obtained by identifying the vertices in each bag, deleting loops, and replacing parallel edges by a single edge. Observe that a graph G has a decomposition D of spread 1 if and only if G has a partition whose pattern is a subgraph of D .

A *tree-partition* is a partition whose pattern is a forest. The *tree-partition-width*²⁰ of a graph G is the minimum width of a tree-partition of G , and is denoted by $\text{tpw}(G)$. Tree-partitions were independently introduced by Seese [80] and Halin [43], and have since been investigated by a number of authors [14, 13, 28, 29, 35, 34, 94].

A graph with bounded degree has bounded tree-partition-width if and only if it has bounded tree-width [29]. In particular, Seese [80] proved the lower bound,

$$2 \cdot \text{tpw}(G) \geq \text{tw}(G) + 1,$$

which is tight for even complete graphs. The best known upper bound is

$$(5) \quad \text{tpw}(G) \leq \frac{5}{2}(\text{tw}(G) + 1) \left(\frac{7}{2} \Delta(G) - 1\right),$$

which was obtained by the first author [95] using a minor improvement to a similar result by an anonymous referee of the paper by Ding and Oporowski [28]. See [6, 24, 25, 30, 31, 56] for other results related to tree-width and graph partitions.

Here we consider more general types of partitions. A partition is *planar* if its pattern is planar. A relationship between planar partitions and rectilinear drawings is described in the following lemma.²¹

²⁰Tree-partition-width has also been called *strong tree-width* [80, 14].

²¹Note that Lemma 8.1 bounds the number of crossings per edge; see [61] for related results.

Lemma 8.1. *Every graph G with a planar partition of width p has a rectilinear drawing in which each edge crosses at most $2 \Delta(G) (p - 1)$ other edges. Hence*

$$\overline{\text{cr}}(G) \leq (p - 1) \Delta(G) \|G\|.$$

Proof. Apply the construction from Lemma 4.1 with $s(v) = 1$ for every vertex v . We obtain a rectilinear drawing of G . Consider an edge vw of G . Say v is in bag X , and w is in bag Y . Then vw is drawn inside $D_e(XY)$. Thus, if two edges e_1 and e_2 of G cross, then an endpoint of e_1 and an endpoint of e_2 are in a common bag, and e_1 and e_2 have no endpoint in common. Thus each edge of G crosses at most $2 \Delta(G) (p - 1)$ other edges, and $\overline{\text{cr}}(G) \leq \frac{1}{2} \sum_e 2 \Delta(G) (p - 1) = \Delta(G) (p - 1) \|G\|$. \square

A graph is *outerplanar* if it has a plane drawing with all the vertices on the outerface. Obviously, $\text{cr}^*(G) = 0$ if and only if G is outerplanar. A partition is *outerplanar* if its pattern is outerplanar.

Lemma 8.2. *Every graph G with an outerplanar partition of width p has a convex drawing in which each edge crosses at most $2 \Delta(G) (p - 1)$ other edges. Hence*

$$\text{cr}^*(G) \leq (p - 1) \Delta(G) \|G\|.$$

Proof. Apply the proof of Lemma 8.1 starting from a plane convex drawing of the pattern. \square

Since every forest is outerplanar, every graph G has an outerplanar partition of width $\text{tpw}(G)$. Thus Lemma 8.2 and Equation (5) imply the following quantitative version of Theorem 1.3.

Corollary 8.3. *Every graph G has a convex drawing in which each edge crosses less than*

$$5 \Delta(G) (\text{tw}(G) + 1) (7 \Delta(G) - 1)$$

other edges. Hence

$$\text{cr}^*(G) < \frac{17}{2} (\text{tw}(G) + 1) \Delta(G)^2 \|G\| < \frac{17}{2} \text{tw}(G) (\text{tw}(G) + 1) \Delta(G)^2 |G|. \quad \square$$

Alon et al. [7] proved that every H -minor free graph G has tree-width at most $c(H) \sqrt{|G|}$ for some constant $c(H)$; also see [42, 23]. Thus Corollary 8.3 implies:

Corollary 8.4. *For every graph H there is a constant $c = c(H)$, such that every graph G with no H -minor has a convex drawing in which each edge crosses less than $c \Delta(G)^2 \sqrt{|G|}$ other edges. Hence*

$$\text{cr}^*(G) < c \Delta(G)^2 \sqrt{|G|} \|G\| < c \Delta(G)^2 |G|^{3/2}. \quad \square$$

Note the following result which is converse to Corollary 8.3.

Proposition 8.5. *Suppose that a graph G has a convex drawing such that whenever two edges e and f cross, e or f crosses at most k edges. Then G has tree-width $\text{tw}(G) \leq 3k + 11$.*

Proof. First we construct a strong planar decomposition D of G (in a similar way to the proof of Lemma 4.2). Replace each vertex v of G by the bag $\{v\}$ in D . Orient each edge of G . Replace each crossing between arcs (v, w) and (x, y) of G by the bag $\{v, x\}$ in D . For each arc (v, w) of G , for some vertex x of G , there is an edge $\{v, x\}\{w\}$ in D ; replace this edge by the path $\{v, x\}\{v, w\}\{w\}$. Thus D is a

strong planar decomposition of G with width 2. Observe that the distance between each bag in D and some bag $\{v\}$ on the outerface is at most $\lfloor \frac{k}{2} \rfloor + 1$. Thus D is $(\lfloor \frac{k}{2} \rfloor + 2)$ -outerplanar.²² Bodlaender [11] proved that every d -outerplanar graph has tree-width at most $3d - 1$. Thus D has tree-width at most $3\lfloor \frac{k}{2} \rfloor + 5$. That is, some tree T is a strong decomposition of D with width at most $3\lfloor \frac{k}{2} \rfloor + 6$. By Lemma 3.4 with $p = 2$, G has a strong decomposition isomorphic to T with width at most $6\lfloor \frac{k}{2} \rfloor + 12$. That is, G has tree-width at most $6\lfloor \frac{k}{2} \rfloor + 11$. \square

9. $K_{3,3}$ -minor-free graphs

In this section we prove Theorem 1.2, which gives an upper bound on the rectilinear crossing number of $K_{3,3}$ -minor-free graphs. The proof employs the following characterisation by Wagner [93].

Lemma 9.1 ([93]). *A graph G is $K_{3,3}$ -minor-free if and only if G can be obtained by (≤ 2) -sums from planar graphs and K_5 .*

Lemma 9.2. *Let G be a $K_{3,3}$ -minor-free graph. For every edge e of G , there is a matching M in G with the following properties:*

- $|M| \leq \frac{1}{3}(|G| - 2)$.
- Each edge in M is disjoint from e .
- Contracting M gives a planar graph.

Proof. If G is planar, then the lemma is satisfied with $M = \emptyset$. Suppose that $G = K_5$. Let vw be an edge of G that is disjoint from e . Let $M := \{vw\}$. Then $|M| = 1 = \frac{1}{3}(|G| - 2)$. The graph obtained by contracting vw is K_4 , which is planar.

Now assume that G is not planar and not K_5 . By Lemma 9.1, G is a (≤ 2) -sum of two smaller $K_{3,3}$ -minor-free graphs G_1 and G_2 . Then $e \in E(G_1)$ or $e \in E(G_2)$. Without loss of generality, $e \in E(G_1)$. If the join set of the (≤ 2) -sum is an edge, then let vw be this edge. Otherwise, let vw be any edge of G_2 .

By induction, G_1 has a matching M_1 with the claimed properties (with respect to the edge e), and G_2 has a matching M_2 with the claimed properties (with respect to the edge vw). In particular, every edge in M_2 is disjoint from vw . Thus $M := M_1 \cup M_2$ is a matching of G (even if $vw \in M_1$). Moreover, every edge in M is disjoint from e . We have $|M| = |M_1| + |M_2| \leq \frac{1}{3}(|G_1| - 2) + \frac{1}{3}(|G_2| - 2) = \frac{1}{3}(|G_1| + |G_2| - 4) \leq \frac{1}{3}(|G| - 2)$.

Let H_i be the planar graph obtained by contracting M_i in G_i . Let H be the graph obtained by contracting M in G . Then H is a (≤ 2) -sum of H_1 and H_2 . Thus H is planar. \square

Corollary 9.3. *Every $K_{3,3}$ -minor-free graph G has a planar partition with width 2 and at most $\frac{1}{3}(|G| - 2)$ bags of cardinality 2.* \square

It follows from Euler’s Formula and Lemma 9.1 that every $K_{3,3}$ -minor-free graph G has at most $3|G| - 5$ edges. Thus Corollary 9.3 and Lemma 8.1 imply the following quantitative version of Theorem 1.2.

²²An outerplanar graph is called 1-outerplanar. A plane graph is k -outerplanar if the graph obtained by deleting the vertices on the outerface is $(k - 1)$ -outerplanar.

Corollary 9.4. *Every $K_{3,3}$ -minor-free graph G has a rectilinear drawing in which each edge crosses at most $2\Delta(G)$ other edges. Hence*

$$\overline{cr}(G) \leq \Delta(G) \|G\| \leq \Delta(G) (3|G| - 5). \quad \square$$

Acknowledgements

Thanks to János Pach and Géza Tóth for explaining their proofs in reference [63]. Thanks to Jaroslav Nešetřil, Patrice Ossona De Mendez, and Vida Dujmović for helpful comments.

An extended abstract of this paper was published in the *Proceedings of the 14th International Symposium on Graph Drawing (GD '06), Lecture Notes in Computer Science*, vol. 4372, pp. 150–161, Springer, 2007.

References

- [1] ADAMSSON, JAY; RICHTER, R. BRUCE. Arrangements, circular arrangements and the crossing number of $C_7 \times C_n$. *J. Combin. Theory Ser. B* **90** (2004), no. 1, 21–39. [MR2041316](#) (2005f:05040), [Zbl 1033.05026](#).
- [2] AGNEW, MELANIE J. It's in the bag: Plane decompositions as tools for approximation. Master's thesis, Department of Computer Science, Rochester Institute of Technology, Rochester, New York, U.S.A., 2005.
- [3] AICHHOLZER, OSWIN; AURENHAMMER, FRANZ; KRASSER, HANNES. On the crossing number of complete graphs. *Computing* **76** (2006), no. 1, 165–176. [MR2174677](#) (2006f:05047), [Zbl 1082.05027](#).
- [4] AIGNER, MARTIN; ZIEGLER, GÜNTER M. *Proofs from The Book*. Springer, third edn., 2004. [MR2014872](#) (2004h:00002), [Zbl 1098.00001](#).
- [5] AJTAI, MIKLÓS; CHVÁTAL, VAŠEK; NEWBORN, MONROE M.; SZEMERÉDI, ENDRE. Crossing-free subgraphs. *Theory and practice of combinatorics*, 9–12, North-Holland Math. Stud., 60, North-Holland, Amsterdam, 1982. [MR806962](#) (86k:05059), [Zbl 0502.05021](#).
- [6] ALON, NOGA; DING, GUOLI; OPOROWSKI, BOGDAN; VERTIGAN, DIRK. Partitioning into graphs with only small components. *J. Combin. Theory Ser. B* **87** (2003), no. 2, 231–243. [MR1957474](#) (2004c:05146), [Zbl 1023.05045](#).
- [7] ALON, NOGA; SEYMOUR, PAUL D.; THOMAS, ROBIN. A separator theorem for nonplanar graphs. *J. Amer. Math. Soc.* **3** (1990), no. 4, 801–808. [MR1065053](#) (91h:05076), [Zbl 0747.05051](#).
- [8] BHATT, SANDEEP N.; LEIGHTON, F. THOMSON. A framework for solving VLSI graph layout problems. *J. Comput. System Sci.* **28** (1984), no. 2, 300–343. [MR0760549](#) (86g:68139).
- [9] BIEDL, THERESE; DEMAINE, ERIK D.; DUNCAN, CHRISTIAN A.; FLEISCHER, RUDOLF; KOBOUROV, STEPHEN G. Tight bounds on maximal and maximum matchings. *Discrete Math.* **285** (2004), no. 1–3, 7–15. [MR2074836](#) (2005b:05179), [Zbl 1044.05056](#).
- [10] BIENSTOCK, DANIEL; DEAN, NATHANIEL. Bounds for rectilinear crossing numbers. *J. Graph Theory* **17** (1993), no. 3, 333–348. [MR1220994](#) (94e:05134), [Zbl 0777.05049](#).
- [11] BODLAENDER, HANS L. Planar graphs with bounded treewidth. Tech. Rep. RUU-CS-88-14, Department of Computer Science, University of Utrecht, Netherlands, 1988.
- [12] BODLAENDER, HANS L. A partial k -arboretum of graphs with bounded treewidth. *Theoret. Comput. Sci.* **209** (1998), no. 1-2, 1–45. [MR1647486](#) (99k:05146), [Zbl 0912.68148](#).
- [13] BODLAENDER, HANS L. A note on domino treewidth. *Discrete Math. Theor. Comput. Sci.* **3** (1999), no. 4, 141–150. [MR1691565](#) (2000b:05078), [Zbl 0912.68148](#).
- [14] BODLAENDER, HANS L.; ENGELFRIET, JOOST. Domino treewidth. *J. Algorithms* **24** (1997), no. 1, 94–123. [MR1453952](#) (98m:68200), [Zbl 0882.68106](#).
- [15] BOKAL, DRAGO. On the crossing numbers of cartesian products with paths. *J. Combin. Theory Ser. B* **97** (2007), no. 3, 381–384.
- [16] BOKAL, DRAGO; CZABARKA, ÉVA; SZÉKELY, LÁSZLÓ A.; VRTO, IMRICH. Graph minors and the crossing number of graphs. *Electron. Notes Discrete Math.* **28** (2007), 169–175.

- [17] BOKAL, DRAGO; FIJAVŽ, GAŠPER; MOHAR, BOJAN. The minor crossing number. *SIAM J. Discrete Math.* **20** (2006), no. 2, 344–356. [MR2257266](#).
- [18] BOKAL, DRAGO; FIJAVŽ, GAŠPER; WOOD, DAVID R. The minor crossing number of graphs with an excluded minor (2006). [arXiv:math/0609707](#).
- [19] BÖRÖCZKY, KÁROLY; PACH, JÁNOS; TÓTH, GÉZA. Planar crossing numbers of graphs embeddable in another surface. *Internat. J. Found. Comput. Sci.* **17** (2006), no. 5, 1005–1015. [MR2270948](#), [Zbl 1100.68075](#).
- [20] BRODSKY, ALEX; DUROCHER, STEPHANE; GETHNER, ELLEN. Toward the rectilinear crossing number of K_n : new drawings, upper bounds, and asymptotics. *Discrete Math.* **262** (2003), no. 1–3, 59–77. [MR1951377](#) (2003i:05039), [Zbl 1013.05027](#).
- [21] CLARK, LANE H.; ENTRINGER, ROGER C. The bisection width of cubic graphs. *Bull. Austral. Math. Soc.* **39** (1989), no. 3, 389–396. [MR0995136](#) (90m:05072), [Zbl 0681.05040](#).
- [22] CZABARKA, ÉVA; SÝKORA, ONDREJ; SZÉKELY, LÁSZLÓ A.; VRT’O, IMRICH. Outerplanar crossing numbers, the circular arrangement problem and isoperimetric functions. *Electron. J. Combin.* **11** (2004), no. 1, R81. [MR2114185](#) (2006b:05040), [Zbl 1080.05022](#).
- [23] DEMAINE, ERIC D.; HAJIAGHAYI, MOHAMMADTAGHI. Graphs excluding a fixed minor have grids as large as treewidth, with combinatorial and algorithmic applications through bidimensionality. *Proc. 16th Annual ACM-SIAM Symp. on Discrete Algorithms* (SODA ’05), 682–689. *ACM*, 2005.
- [24] DEMAINE, ERIK D.; HAJIAGHAYI, MOHAMMADTAGHI; KAWARABAYASHI, KEN-ICHI. Algorithmic graph minor theory: Decomposition, approximation, and coloring. *Proc. 46th Annual IEEE Symposium on Foundations of Computer Science* (FOCS ’05), 637–646. *IEEE*, 2005.
- [25] DEVOS, MATT; DING, GUOLI; OPOROWSKI, BOGDAN; SANDERS, DANIEL P.; REED, BRUCE; SEYMOUR, PAUL; VERTIGAN, DIRK. Excluding any graph as a minor allows a low tree-width 2-coloring. *J. Combin. Theory Ser. B* **91** (2004), no. 1, 25–41. [MR2047529](#) (2004m:05247), [Zbl 1042.05036](#).
- [26] DÍAZ, JOSEP; DO, NORMAN; SERNA, MARIA J.; WORMALD, NICHOLAS C. Bounds on the max and min bisection of random cubic and random 4-regular graphs. *Theoret. Comput. Sci.* **307** (2003), no. 3, 531–547. [MR2021746](#) (2005h:05184), [Zbl 1070.68119](#).
- [27] DIESTEL, REINHARD; KÜHN, DANIELA. Graph minor hierarchies. *Discrete Appl. Math.* **145** (2005), no. 2, 167–182. [MR2113139](#) (2005i:05180), [Zbl 1055.05140](#).
- [28] DING, GUOLI; OPOROWSKI, BOGDAN. Some results on tree decomposition of graphs. *J. Graph Theory* **20** (1995), no. 4, 481–499. [MR1358539](#) (96m:05053), [Zbl 0837.05044](#).
- [29] DING, GUOLI; OPOROWSKI, BOGDAN. On tree-partitions of graphs. *Discrete Math.* **149** (1996), no. 1–3, 45–58. [MR1375097](#) (97a:05168), [Zbl 0844.05078](#).
- [30] DING, GUOLI; OPOROWSKI, BOGDAN; SANDERS, DANIEL P.; VERTIGAN, DIRK. Partitioning graphs of bounded tree-width. *Combinatorica* **18** (1998), no. 1, 1–12. [MR1645638](#) (99i:05162), [Zbl 0924.05022](#).
- [31] DING, GUOLI; OPOROWSKI, BOGDAN; SANDERS, DANIEL P.; VERTIGAN, DIRK. Surfaces, tree-width, clique-minors, and partitions. *J. Combin. Theory Ser. B* **79** (2000), no. 2, 221–246. [MR1769192](#) (2001i:05059), [Zbl 1029.05041](#).
- [32] DJIDJEV, HRISTO N.; VRT’O, IMRICH. Crossing numbers and cutwidths. *J. Graph Algorithms Appl.* **7** (2003), no. 3, 245–251. [MR2112230](#) (2005h:05055), [Zbl 1066.05054](#).
- [33] DJIDJEV, HRISTO N.; VRŤO, IMRICH. Planar crossing numbers of genus g graphs. *Proc. 33rd International Colloquium on Automata, Languages and Programming* (ICALP ’06) (BUGLIESI, MICHELE; PRENEEL, BART; SASSONE, VLADIMIRO; WEGENER, INGO, eds.), Lecture Notes in Comput. Sci., 4051, 419–430. Springer, 2006.
- [34] DUJMOVIĆ, VIDA; MORIN, PAT; WOOD, DAVID R. Layout of graphs with bounded tree-width. *SIAM J. Comput.* **34** (2005), no. 3, 553–579. [MR2137079](#) (2006a:05103), [Zbl 1069.05055](#).
- [35] EDENBRANDT, ANDERS. Quotient tree partitioning of undirected graphs. *BIT* **26** (1986), no. 2, 148–155. [MR0840323](#) (87g:05077), [Zbl 0591.68070](#).
- [36] ERDŐS, PAUL; GUY, RICHARD K. Crossing number problems. *Amer. Math. Monthly* **80** (1973), 52–58. [MR0382006](#), [Zbl 0264.05109](#).
- [37] FÁRY, ISTVÁN. On straight line representation of planar graphs. *Acta Univ. Szeged. Sect. Sci. Math.* **11** (1948), 229–233. [MR0026311](#) (10,136f), [Zbl 0030.17902](#).

- [38] GARCIA-MORENO, ENRIQUE; SALAZAR, GELASIO. Bounding the crossing number of a graph in terms of the crossing number of a minor with small maximum degree. *J. Graph Theory* **36** (2001), no. 3, 168–173. [MR1814533](#) (2001m:05093), [Zbl 0979.05033](#).
- [39] GAREY, MICHAEL R.; JOHNSON, DAVID S. Crossing number is NP-complete. *SIAM J. Algebraic Discrete Methods* **4** (1983), no. 3, 312–316. [MR0711340](#) (85a:68070), [Zbl 0536.05016](#).
- [40] GEELLEN, JAMES F.; RICHTER, R. BRUCE; SALAZAR, GELASIO. Embedding grids in surfaces. *European J. Combin.* **25** (2004), no. 6, 785–792. [MR2079898](#) (2005c:05057), [Zbl 1050.05035](#).
- [41] GLEBSKY, LEV YU.; SALAZAR, GELASIO. The crossing number of $C_m \times C_n$ is as conjectured for $n \geq m + 1$. *J. Graph Theory* **47** (2004), no. 1, 53–72. [MR2079290](#) (2005c:05058), [Zbl 1053.05032](#).
- [42] GROHE, MARTIN. Local tree-width, excluded minors, and approximation algorithms. *Combinatorica* **23** (2003), no. 4, 613–632. [MR2046826](#) (2004m:05248), [Zbl 1089.05503](#).
- [43] HALIN, RUDOLF. Tree-partitions of infinite graphs. *Discrete Math.* **97** (1991), 203–217. [MR1140802](#) (93b:05044), [Zbl 0763.05027](#).
- [44] HLINĚNÝ, PETR. Crossing-number critical graphs have bounded path-width. *J. Combin. Theory Ser. B* **88** (2003), no. 2, 347–367. [MR1983364](#) (2004c:05060), [Zbl 1021.05028](#).
- [45] HLINĚNÝ, PETR. Crossing number is hard for cubic graphs. *J. Combin. Theory Ser. B* **96** (2006), no. 4, 455–471. [MR2232384](#) (2007c:05056), [Zbl 1092.05016](#).
- [46] KAWARABAYASHI, KEN-ICHI; MOHAR, BOJAN. Some recent progress and applications in graph minor theory. *Graphs Combin.* **23** (2007), no. 1, 1–46.
- [47] KOSTOCHKA, ALEXANDR V. The minimum Hadwiger number for graphs with a given mean degree of vertices. *Metody Diskret. Analiz.* **38** (1982), 37–58. [MR0713722](#) (85c:05018), [Zbl 0544.05037](#).
- [48] LEAÑOS, JESÚS; SALAZAR, GELASIO. On the additivity of crossing numbers of graphs, 2006. <http://www.ifisica.uaslp.mx/~gsalazar/RESEARCH/additivity.pdf>.
- [49] LEIGHTON, F. THOMSON. Complexity issues in VLSI. *MIT Press*, 1983. ISBN 0262121042.
- [50] LEIGHTON, F. THOMSON. New lower bound techniques for VLSI. *Math. Systems Theory* **17** (1984), no. 1, 47–70. [MR0738751](#) (85d:68040), [Zbl 0488.94048](#).
- [51] LOVÁSZ, LÁSZLÓ; VESZTERGOMBI, KATALIN; WAGNER, ULI; WELZL, EMO. Convex quadrilaterals and k -sets. In PACH [57], 139–148. [MR2065247](#) (2004m:05004), [Zbl 1052.05004](#).
- [52] MOHAR, BOJAN; THOMASSEN, CARSTEN. Graphs on surfaces. *Johns Hopkins University Press, Baltimore, U.S.A.*, 2001. [MR1844449](#) (2002e:05050), [Zbl 0979.05002](#).
- [53] MONTARON, BERNARD. An improvement of the crossing number bound. *J. Graph Theory* **50** (2005), no. 1, 43–54. [MR2157537](#) (2006b:05047), [Zbl 1073.05020](#).
- [54] NAHAS, NAGI H. On the crossing number of $K_{m,n}$. *Electron. J. Combin.* **10** (2003), N8. [MR2014536](#) (2004i:05039), [Zbl 1023.05039](#).
- [55] NEGAMI, SEIYA. Crossing numbers of graph embedding pairs on closed surfaces. *J. Graph Theory* **36** (2001), no. 1, 8–23. [MR1803630](#) (2001m:05097), [Zbl 0971.05037](#).
- [56] NEŠETŘIL, JAROSLAV; OSSONA DE MENDEZ, PATRICE. Grad and classes with bounded expansion I. Decompositions. *European J. Combin.* (in press). <http://dx.doi.org/10.1016/j.ejc.2006.07.013>.
- [57] PACH, JÁNOS, ed. Towards a theory of geometric graphs, Contemporary Mathematics, 342. *Amer. Math. Soc., Providence*, 2004. [MR2065247](#) (2004m:05004), [Zbl 1052.05004](#).
- [58] PACH, JÁNOS; RADOIČIĆ, RADOŠ; TARDOS, GÁBOR; TÓTH, GÉZA. Improving the crossing lemma by finding more crossings in sparse graphs. *Discrete Comput. Geom.* **36** (2006), no. 4, 527–552. [MR2267545](#), [Zbl 1104.05022](#).
- [59] PACH, JÁNOS; SHAHROKHI, FARHAD; SZEGEDY, MARIO. Applications of the crossing number. *Algorithmica* **16** (1996), no. 1, 111–117. [MR1394496](#) (97h:05054), [Zbl 0851.68088](#).
- [60] PACH, JÁNOS; SHARIR, MICHA. On the number of incidences between points and curves. *Combin. Probab. Comput.* **7** (1998), no. 1, 121–127. [MR1611057](#) (99b:52037), [Zbl 0901.52016](#).
- [61] PACH, JÁNOS; TÓTH, GÉZA. Graphs drawn with few crossings per edge. *Combinatorica* **17** (1997), no. 3, 427–439. [MR1606052](#) (99b:05043), [Zbl 0902.05017](#).
- [62] PACH, JÁNOS; TÓTH, GÉZA. Which crossing number is it anyway? *J. Combin. Theory Ser. B* **80** (2000), no. 2, 225–246. [MR1794693](#) (2001h:05034), [Zbl 1023.05042](#).
- [63] PACH, JÁNOS; TÓTH, GÉZA. Crossing number of toroidal graphs. *Proc. 13th International Symp. on Graph Drawing (GD '05)* (HEALY, PATRICK; NIKOLOV, NIKOLA S., eds.), Lecture Notes in Comput. Sci., 3843, 334–342. *Springer, Berlin*, 2006. [MR2244523](#).

- [64] PELSMAJER, MICHAEL J.; SCHAEFER, MARCUS; ŠTEFANKOVIČ, DANIEL. Crossing number of graphs with rotation systems. Tech. Rep. 05-017, School of Computer Science, Telecommunications and Information Systems, DePaul University, Chicago, U.S.A., 2005.
- [65] PURCHASE, HELEN. Which aesthetic has the greatest effect on human understanding? *Proc. 5th International Symp. on Graph Drawing (GD '97)* (DI BATTISTA, GIUSEPPE, ed.), Lecture Notes in Comput. Sci., 1353, 248–261. Springer, 1997.
- [66] PURCHASE, HELEN C. Performance of layout algorithms: Comprehension, not computation. *J. Visual Languages and Computing* **9** (1998), 647–657.
- [67] PURCHASE, HELEN C.; COHEN, ROBERT F.; JAMES, MURRAY I. An experimental study of the basis for graph drawing algorithms. *ACM J. Experimental Algorithmics* **2** (1997), no. 4. <http://www.jea.acm.org/1997/PurchaseDrawing/>.
- [68] REED, BRUCE A. Algorithmic aspects of tree width. *Recent Advances in Algorithms and Combinatorics* (REED, BRUCE A.; SALES, CLÁUDIA L., eds.), 85–107. CMS Books Math./Ouvrages Math. SMC, 11, Springer, New York, 2003. [MR1952984](#) (2003m:05199), [Zbl 1035.05090](#).
- [69] RICHTER, R. BRUCE; ŠIRÁŇ, JOZEF. The crossing number of $K_{3,n}$ in a surface. *J. Graph Theory* **21** (1996), no. 1, 51–54. [MR1363689](#) (96i:05054), [Zbl 0838.05033](#).
- [70] RICHTER, R. BRUCE; THOMASSEN, CARSTEN. Intersections of curve systems and the crossing number of $C_5 \times C_5$. *Discrete Comput. Geom.* **13** (1995), no. 2, 149–159. [MR1314959](#) (95j:05081), [Zbl 0820.05015](#).
- [71] RICHTER, R. BRUCE; THOMASSEN, CARSTEN. Relations between crossing numbers of complete and complete bipartite graphs. *Amer. Math. Monthly* **104** (1997), no. 2, 131–137. [MR1437414](#) (98f:05054), [Zbl 0872.05010](#).
- [72] RINGEL, GERHARD. Das Geschlecht des vollständigen paaren Graphen. *Abh. Math. Sem. Univ. Hamburg* **28** (1965), 139–150. [MR0189012](#), [Zbl 0132.21203](#).
- [73] RINGEL, GERHARD. *Map color theorem*. Springer-Verlag, 1974. [MR0349461](#), [Zbl 0287.05102](#).
- [74] RISKIN, ADRIAN. On the outerplanar crossing numbers of $K_{m,n}$. *Bull. Inst. Combin. Appl.* **39** (2003), 16–20. [MR1999850](#), [Zbl 1048.05033](#).
- [75] ROBERTSON, NEIL; SANDERS, DANIEL P.; SEYMOUR, PAUL D.; THOMAS, ROBIN. The four-colour theorem. *J. Combin. Theory Ser. B* **70** (1997), no. 1, 2–44. [MR1441258](#) (98c:05065), [Zbl 0883.05056](#).
- [76] ROBERTSON, NEIL; SEYMOUR, PAUL D. Graph minors. II. Algorithmic aspects of tree-width. *J. Algorithms* **7** (1986), no. 3, 309–322. [MR0855559](#) (88c:05053), [Zbl 0611.05017](#).
- [77] ROBERTSON, NEIL; SEYMOUR, PAUL D. Graph minors. XVI. Excluding a non-planar graph. *J. Combin. Theory Ser. B* **89** (2003), no. 1, 43–76. [MR1999736](#) (2004f:05168), [Zbl 1023.05040](#).
- [78] ROBERTSON, NEIL; SEYMOUR, PAUL D. Graph minors. XX. Wagner’s conjecture. *J. Combin. Theory Ser. B* **92** (2004), no. 2, 325–357. [MR2099147](#) (2005m:05204), [Zbl 1061.05088](#).
- [79] SCHEINERMAN, EDWARD R.; WILF, HERBERT S. The rectilinear crossing number of a complete graph and Sylvester’s “four point problem” of geometric probability. *Amer. Math. Monthly* **101** (1994), no. 10, 939–943. [MR1304316](#) (95k:52006), [Zbl 0834.05022](#).
- [80] SEESE, DETLEF. Tree-partite graphs and the complexity of algorithms. *Proc. International Conf. on Fundamentals of Computation Theory* (BUDACH, LOTHAR, ed.), Lecture Notes in Comput. Sci., 199, 412–421. Springer, Berlin, 1985. [MR0821258](#), [Zbl 0574.68036](#).
- [81] SEESE, DETLEF G.; WESSEL, WALTER. Grids and their minors. *J. Combin. Theory Ser. B* **47** (1989), no. 3, 349–360. [MR1026069](#) (90j:05127), [Zbl 0636.05023](#).
- [82] SHAHROKHI, FARHAD; SÝKORA, ONDREJ; SZÉKELY, LÁSZLÓ A.; VRŤO, IMRICH. The crossing number of a graph on a compact 2-manifold. *Adv. Math.* **123** (1996), no. 2, 105–119. [MR1420482](#) (97h:05055), [Zbl 0865.05036](#).
- [83] SHAHROKHI, FARHAD; SÝKORA, ONDREJ; SZÉKELY, LASZLO A.; VRŤO, IMRICH. The gap between crossing numbers and convex crossing numbers. In PACH [57], 249–258. [MR2065268](#) (2005c:05064), [Zbl 1062.05050](#).
- [84] SHAHROKHI, FARHAD; SZÉKELY, LÁSZLÓ A. On canonical concurrent flows, crossing number and graph expansion. *Combin. Probab. Comput.* **3** (1994), no. 4, 523–543. [MR1314073](#) (96i:05090), [Zbl 0817.05022](#).
- [85] SHAHROKHI, FARHAD; SZÉKELY, ONDREJ, LÁSZLÓ A. SÝKORA; VRŤO, IMRICH. Drawings of graphs on surfaces with few crossings. *Algorithmica* **16** (1996), no. 1, 118–131. [MR1394497](#) (97h:05056), [Zbl 0854.68074](#).

- [86] SZÉKELY, LÁSZLÓ A. Crossing numbers and hard Erdős problems in discrete geometry. *Combin. Probab. Comput.* **6** (1997), no. 3, 353–358. [MR1464571](#) (98h:52030), [Zbl 0882.52007](#).
- [87] SZÉKELY, LÁSZLÓ A. A successful concept for measuring non-planarity of graphs: the crossing number. *Discrete Math.* **276** (2004), no. 1–3, 331–352. [MR2046646](#) (2005a:05073), [Zbl 1035.05034](#).
- [88] TAIT, PETER GUTHRIE. Note on a theorem in geometry of position. *Trans. Roy. Soc. Edinburgh* **29** (1880), 657–660.
- [89] THOMASON, ANDREW. An extremal function for contractions of graphs. *Math. Proc. Cambridge Philos. Soc.* **95** (1984), no. 2, 261–265. [MR0735367](#) (85i:05141), [Zbl 0551.05047](#).
- [90] THOMASON, ANDREW. The extremal function for complete minors. *J. Combin. Theory Ser. B* **81** (2001), no. 2, 318–338. [MR1814910](#) (2002i:05100), [Zbl 1024.05083](#).
- [91] VRŤO, IMRICH. Crossing numbers of graphs: A bibliography (2007). <ftp://ftp.if.savba.sk/pub/imrich/crobib.pdf>.
- [92] WAGNER, KLAUS. Bemerkung zum Vierfarbenproblem. *Jber. Deutsch. Math.-Verein.* **46** (1936), 26–32.
- [93] WAGNER, KLAUS. Über eine Eigenschaft der ebene Komplexe. *Math. Ann.* **114** (1937), 570–590. [MR1513158](#), [Zbl 0017.19005](#).
- [94] WOOD, DAVID R. Vertex partitions of chordal graphs. *J. Graph Theory* **53** (2006), no. 2, 167–172. [MR2255760](#).
- [95] WOOD, DAVID R. A note on tree-partition-width (2007). [arXiv:math/0602507](#).
- [96] WOOD, DAVID R.; TELLE, JAN ARNE. Planar decompositions and the crossing number of graphs with an excluded minor (2006). [arXiv:math/0604467](#).

DEPARTAMENT DE MATEMÀTICA APLICADA II, UNIVERSITAT POLITÈCNICA DE CATALUNYA,
BARCELONA, SPAIN
david.wood@upc.es

DEPARTMENT OF INFORMATICS, THE UNIVERSITY OF BERGEN, BERGEN, NORWAY
Jan.Arne.Telle@ii.uib.no

This paper is available via <http://nyjm.albany.edu/j/2007/13-8.html>.